

CELL TOWERS NEAR SCHOOLS

SCHOOL CELL TOWER SETBACKS

Many communities have policies, ordinances or zoning that ensures cellular antennas are restricted to a specific minimum distance from schools. Hempstead, New York requires a special use permit for cell towers near schools.

Examples of cell tower/4G/5G small cell setbacks/preferred placements for schools:

- Palo Alto, California: 1,500 feet
- Copake, New York: 1,500 feet
- Los Altos, California: 500 feet
- Walnut City, California: 1,500 feet
- Bar Harbor, Maine: 1,500 feet
- Sallisaw, Oklahoma: 1,500 feet
- Shelbourne, Massachusetts: 1,500 feet
- Stockbridge, Massachusetts: 1,500 feet
- San Diego County California: 1,000 feet
- Encinitas California: 500 feet
- Scarsdale New York: 500 feet
- Ithaca, New York: 250 feet

CELL TOWERS REMOVED FROM SCHOOL GROUNDS

- Milpitas California: School Board asked Crown Castle and T-Mobile to relocate the cell tower to remote location.
- Ripon California: Sprint moved the cell tower at elementary after students and staff developed cancer and parents argued children should not be guinea pigs.
- Alameda California cancelled cell tower contracts.
- Dekalb County Georgia dropped school tower plan.

SCHOOL BOARDS

- Palo Alto Unified School District Cell Tower Resolution supports the City 1,500 setback and opposes cell tower "on or in close proximity to schools to ensure individuals, especially children, are protected from the potential negative effects associated with radiation exposure."
- West Linn-Wilsonville Oregon School Board prohibits cell towers on school property.
- Vancouver School Boards Resolution: 1,000 feet
- Greenbelt Maryland Council opposes school towers.

DID YOU KNOW?

- The International Association of Firefighters passed a Resolution opposing cell towers on its stations in 2004 after a study found neurological damage in firefighters with antennas on their fire stations.

LOS ANGELES UNIFIED CA SCHOOL DISTRICT

- 3 resolutions opposing cell towers on school property.
- The District Office of Health and Safety developed a "cautionary level" for radiofrequency radiation 10,000 times lower than FCC regulations because, "it is believed that a more conservative level is necessary to protect children, who represent a potentially vulnerable and sensitive population."

SCHOOL BOARDS THAT REVERSED COURSE

- Montgomery County Maryland Schools has a policy no to allow any more cell towers on elementary schools.
- Prince George's County Maryland School Board decided not to renew a cell tower construction master leasing agreement that had allowed over 70 schools to be marketed as cell tower sites.
- Portland Oregon Schools ended new leases for cell towers.

EXPERT RECOMMENDATIONS

- The New Hampshire State Commission 5G Health and Environment Report recommends a setback of 1640 feet for schools.
- The Collaborative For High Performance Schools (Green building rating program) has LOW EMF Criteria which includes no cell towers on school property.

THE EPA SCHOOL SITING GUIDELINES

Lists exposure to electromagnetic fields and the fall distance as "potential hazards" from cell towers. The EPA guidelines recommend schools "identify and evaluate cell towers within ~200 feet of prospective school locations."

PUBLISHED RESEARCH

- 500 meter buffer recommended for schools to reduce liability and minimize risk (Pearce 2019)
- A moratorium on 5G pending safety research (Frank 2020)
- A precautionary approach is better suited to State obligations under international human rights law (Roda and Perry 2014)
- Increased cancer deaths near cell antennas (Rodrigues 2021)
- Studies find: DNA Damage (Zothansiana 2017), Diabetes (Meo 2015), Cognitive effects (Meo 2018), sleep problems and headaches (Abdel-Rassoul 2007, Levitt & Lai 2010, Shahbazi-Gahrouei 2013)

5G, CELL TOWERS AND WIRELESS LEGAL & LIABILITY ISSUES



When a new cell tower is proposed, the first question to ask is: "Do you have insurance for damages from long-term exposure to the radiofrequency radiation (RFR)?"

Usually the answer is "No." Why? Insurance companies rank the risk as "HIGH."

5G and Cell Towers Are an Uninsurable Risk

- Insurers rank wireless, cell tower, and 5G RFR non-ionizing electromagnetic field (EMF) radiation as a "high" risk, comparing the issue to lead and asbestos.
- Most insurance plans have "electromagnetic field exclusions" and do not insure for long-term RFR damages.
- Additionally, some insurance plans will not provide a defense for any supervision instruction or recommendation given "or which should have been given" in connection to EMFs.
- Wireless RFR and non-ionizing electromagnetic radiation are defined as a type of "pollution" by wireless companies themselves.
- U.S. mobile operators have been unable to get insurance to cover liabilities related to damages from long-term RFR exposure.
- Wireless companies warn their shareholders of RFR risk but do not warn users of their products, nor do the companies warn the people exposed to emissions from their infrastructure.

Cell Tower Companies Warn Shareholders of Risk From Cell Tower Radiation

Why Don't They Warn Families Living Near Cell Towers?



Verizon 10-K Report

"Our wireless business also faces personal injury and wrongful death lawsuits relating to alleged health effects of wireless phones or radio frequency transmitters. We may incur significant expenses in defending these lawsuits. In addition, we may be required to pay significant awards or settlements."

Crown Castle 10-K Report

"We cannot guarantee that claims relating to radio frequency emissions will not arise in the future or that the results of such studies will not be adverse to us...If a connection between radio frequency emissions and possible negative health effects were established, our operations, costs, or revenues may be materially and adversely affected. We currently do not maintain any significant insurance with respect to these matters."



AT&T 10-K Report

"In the wireless area, we also face current and potential litigation relating to alleged adverse health effects on customers or employees who use such technologies including, for example, wireless devices. We may incur significant expenses defending such suits or government charges and may be required to pay amounts or otherwise change our operations in ways that could materially adversely affect our operations or financial results."



T- MOBILE 10-K Report

"Our business could be adversely affected by findings of product liability for health or safety risks from wireless devices and transmission equipment, as well as by changes to regulations or radio frequency emission standards."



Cell Tower Companies Warn Shareholders of Risk From Cell Tower Radiation

Why Don't They Warn Families Living Near Cell Towers?



AMERICAN TOWER®

NOKIA
CONNECTING PEOPLE



Qualcomm



ERICSSON

American Tower 10-K

"If a scientific study or court decision resulted in a finding that radio frequency emissions pose health risks to consumers, it could negatively impact our tenants and the market for wireless services, which could materially and adversely affect our business, results of operations or financial condition. We do not maintain any significant insurance with respect to these matters."

Nokia 10-K

"Although our products are designed to meet all relevant safety standards and other recommendations and regulatory requirements globally, we cannot guarantee we will not become subject to product liability claims or be held liable for such claims, which could have a material adverse effect on us."

Qualcomm 10-K

"If wireless handsets pose health and safety risks, we may be subject to new regulations, and demand for our products and those of our licensees and customers may decrease."

Ericsson Annual Report

"Any perceived risk or new scientific findings of adverse health effects from mobile communication devices and equipment could adversely affect us through a reduction in sales or through liability claims."

T-Mobile Warns of the Risk of 5G and Lawsuits

The Data on Risk Could Change, Impacting Cash Flow



T-Mobile advertises to the public about going "live" but omits the warnings they give to shareholders regarding 5G, regulatory changes and risk perception.

T-Mobile™

T-Mobile 10-K Report 2/2023

"Negative public perception of, and regulations regarding, the perceived health risks relating to 5G networks could undermine market acceptance of our 5G services" (page 13)

"We, along with equipment manufacturers and other carriers, are subject to current and potential future lawsuits **alleging adverse health effects arising from the use of wireless handsets or from wireless transmission equipment such as cell towers.**"

"In addition, **the FCC has from time to time gathered data regarding wireless device emissions, and its assessment of the risks associated with using wireless devices may evolve based on its findings.** Any of these allegations or changes in risk assessments could result in customers purchasing fewer devices and wireless services, could result in significant legal and regulatory liability, and could have a material adverse effect on our business, reputation, financial condition, cash flows and operating results." (T-Mobile 10-K Report page 21)



A 2000 Ecolog Institute Report commissioned by T-Mobile and DeTeMobil Deutsche Telekom MobilNet recommended an RF exposure limit 1000x lower than the FCC's current power density limit after reviewing the research on biological effects, including impacts to the immune system, central nervous system, hormones, cancer, neurotransmitters and fertility.


This PDF is hyperlinked. For more on legal liability issues go to ehtrust.org

ENVIRONMENTAL HEALTH TRUST | [EHTRUST.ORG](https://ehtrust.org)



5G, CELL TOWERS AND WIRELESS

LEGAL & LIABILITY ISSUES SHAREHOLDER WARNINGS



"Some research has shown biological effects from lower-level "non thermal" exposure and people exposed at lower levels have reported headaches, dizziness, nausea, mood disorders, mental slowing and memory loss."

***Business Insurance White Paper,
The Next Asbestos: Five Emerging Risks
That Could Shift the Liability Landscape***

Insurance Authorities Rate 5G as "High Risk."

5G mobile networks are classified as a "high," "off-the-leash" risk. "Existing concerns regarding potential negative health effects from electromagnetic fields (EMF) are only likely to increase. An uptick in liability claims could be a potential long-term consequence" and "as the biological effects of EMF in general and 5G in particular are still being debated, potential claims for health impairments may come with a long latency."
— Swiss Re Institute (2019)

Insurance Companies Have Electromagnetic Field Exclusions As the Industry Standard

Electromagnetic field exclusions" are clear and common in most insurance companies. It is applied as a market standard. This exclusion serves to exclude cover for illnesses caused by long-term EMF (non-ionizing radiation) exposure." — Complete Markets

"Exclusions: This insurance does not apply to: Bodily injury, personal injury, advertising injury, or property damage arising directly or indirectly out of, resulting from, caused or contributed to by electromagnetic radiation, provided that such loss, cost or expense results from or is contributed to by the hazardous properties of electromagnetic radiation.
— Portland Oregon Public School Insurance (page 30)

Insurance Plans Not Only Exclude EMF Damages, But Some Even Exclude Defending Decision Makers From Actions

"This policy does not apply to and we will not provide a defense for: a. bodily injury... arising out of ... exposure to or contact with electromagnetic radiation... b. costs of abatement .. of EMF" or c. any supervision, instruction, recommendation, warning or advice given or which should have been given in connection with a or b. above."- City of Ann Arbor Michigan Insurance Policy page 14.

Wireless Companies Rank EMF as a Risk with High Impact

"Electro-magnetic signals emitted by mobile devices and base stations may be found to pose health risks, with potential impacts including: changes to national legislation, a reduction in mobile phone usage or litigation."
— Vodaphone 2017 Report ranks EMF as a "Principal Risk with "High" impact.

Wireless Companies Warn Shareholder About Risk But Not People Living Near Their Wireless Infrastructure

Crown Castle says:

"We cannot guarantee that claims relating to radio frequency emissions will not arise in the future or that the results of such studies will not be adverse to us...If a connection between radio frequency emissions and possible negative health effects were established, our operations, costs, or revenues may be materially and adversely affected. We currently do not maintain any significant insurance with respect to these matters."

Wireless Companies Define Pollution in Their Own Policies as Including EMFs, Microwaves and Non-ionizing Radiation.

Verizons Total Mobile Protection Plan says: "Pollution" is defined as "any solid, liquid, gaseous, or thermal irritant or contaminant including smoke, vapor, soot, fumes, acid, alkalis, chemicals, artificially produced electric fields, magnetic field, electromagnetic field, sound waves, microwaves, and all artificially produced ionizing or non-ionizing radiation and/or waste."

You work best when your tech works too.

Total Mobile Protection for Business



Applicable for Business customers outside of New York. New York customers, please see the Total Mobile Protection for Business brochure for New York.


Verizon Total Mobile Protection Plan Defines Non-ionizing Radiation as "Pollution"

16. Pollution

The discharge, dispersal, seepage, migration or escape of pollutants. Pollutants means any solid, liquid, gaseous, or thermal irritant or contaminant including smoke, vapor, soot, fumes, acid, alkalis, chemicals, artificially produced electric fields, magnetic field, electromagnetic field, sound waves, microwaves, and all artificially produced ionizing or non-ionizing radiation and/or waste. Waste includes materials to be recycled, reconditioned or reclaimed.

AT&T, Sprint and T-Mobile also have similar "pollution" definitions and they refuse to cover damages.

Click on image to view the policy.



Insurance Companies Exclude EMF As Industry Standard

THIS ENDORSEMENT CHANGES THE POLICY. PLEASE READ IT CAREFULLY.

ELECTROMAGNETIC RADIATION EXCLUSION

This endorsement modifies insurance provided under the following:

GENERAL LIABILITY COVERAGE PART
PUBLIC RISK GENERAL LIABILITY RETAINED LIMIT COVERAGE FORM
LAW ENFORCEMENT COVERAGE PART
LAW ENFORCEMENT LIABILITY RETAINED LIMIT COVERAGE FORM
PUBLIC OFFICIALS COVERAGE PART
PUBLIC OFFICIALS LIABILITY RETAINED LIMIT COVERAGE FORM
EMPLOYMENT PRACTICES LIABILITY COVERAGE PART
EMPLOYMENT PRACTICES LIABILITY RETAINED LIMIT COVERAGE FORM

The following **Exclusion** is added:

This policy does not apply to and we will not provide a defense for:

- a. "Bodily injury," "property damage", "personal and advertising injury", "employee benefits wrongful acts", "personal injury", "law enforcement wrongful acts", "public officials wrongful acts", "educator's legal wrongful acts", or "employment practices wrongful acts" arising out of, or which result in, the actual, alleged, threatened, perceived, latent, sudden and accidental or incidental exposure to or contact with electromagnetic radiation in any form, from any source.
- b. The costs of abatement or mitigation of:
 - (1) Electromagnetic radiation; or
 - (2) Exposure to electromagnetic radiation.
- c. Any supervision, instruction, recommendation, warning or advice given or which should have been given in connection with a. or b. above.

Electromagnetic radiation includes but is not limited to, magnetic energy, waves, fields or forces generated, produced, transmitted or maintained by the charges, currents, frequencies, energy or forces of electricity that is generated, flowing or otherwise transmitted through or via the medium, methods and equipment designed to generate, produce, distribute, transport or transmit the electrical charges, currents, frequencies, energy or forces.

Click to Download Insurance Terms That Define Pollution as Including Wireless Electromagnetic

16. Pollution



The discharge, dispersal, seepage, migration or escape of pollutants. Pollutants means any solid, liquid, gaseous, or thermal irritant or contaminant including smoke, vapor, soot, fumes, acid, alkalis, chemicals, artificially produced electric fields, magnetic field, electromagnetic field, sound waves, microwaves, and all artificially produced ionizing or non-ionizing radiation and/or waste. Waste includes materials to be recycled, reconditioned or reclaimed.



Sprint
Complete

For Smart Devices
Tier 2-5
Terms and Conditions

Para dispositivos inteligentes
Niveles 2-5
Términos y condiciones



*Welcome to
AT&T Multi-Device
Protection Pack*

Important Information,
including Terms and Conditions


**Premium Handset
Protection**
Insurance
Warranty



**Live your life.
We've got
you covered.**



PARENT TEACHER ASSOCIATIONS OPPOSE CELL TOWERS



CONEJO PTA WANTS CELL TOWER MOVED **Op-ed in Thousand Oaks Acorn Journal**

The California PTA advocates on behalf of children and families. They advocate against electromagnetic field radiation your schools.

The Conejo PTA urges the use of the precautionary principle in making decisions regarding public health this means if something cannot be proven to be safe it is best to avoid exposure. Most people don't realize that the 1996 FCC state standards for safe levels of omission was actually based on a level set by the American national standards institute in 1982. Well this standard has not been changed in 30 years it has usurped all local authority."

"For this reason, Conejo Council PTA made up of 9000 parents and teachers has decided to take action. We're calling on our local leaders to put in place policies that would ensure parents are notified when cell towers are propose near schools and then encourage a buffer zone around schools."

-Kim Huber, legislative chair of the Conejo Council PTA.

NEW YORK STATE PTA **-Adopted TWO Resolutions 2014**

"CELLULAR PHONE TOWERS – 2014 (R-'07, R-'00); Resolved that the New York State Congress of Parents and Teachers, Inc. support legislation that would encourage local communities, including parents and school officials, to regulate the placement of cell towers and cell tower antennas particularly in schools and areas where children congregate,

and be it further Resolved that the New York State PTA support continued research into the long-term effects of radio frequency and microwave frequencies on humans especially as they apply to children, and be it further Resolved that the New York State PTA seek to educate parents and school officials as to the current debate over the placement of cell towers and antennas."

NEELSVILLE MIDDLE SCHOOL PTA (MD)

- Voted to oppose proposed cell tower.
- Hosted parent information session with both the cell tower company and Environmental Health Trust.

HILLSMERE ELEMENTARY SCHOOL PTA (MD)

- Sent letters to the school board in opposition to cell towers near the school.

BRIARLAKE ELEMENTARY (GA)


- Voted to oppose cell tower after board approved towers on schools.

PACIFIC GROVE (CA) PTAs

- Forest Grove Elementary Pacific Grove Middle School and Pacific Grove High School PTAs sent a letter to City Council opposing a high school cell tower.

UNITED STATES OF AMERICA

5G & CELL TOWERS



CALIFORNIA

Numerous CA cities restrict cell antennas near homes with setbacks and strict ordinances including: Los Altos, Petaluma, Mill Valley, Malibu, Santa Barbara, Nevada City, Suisin, Calabasas, San Clemente, Westlake, Sonoma, Sebastopol, San Rafael, Ross Valley, Encinitas, Fairfax, Palo Alto, Walnut City and San Diego County.

As an example of CA ordinances, the Los Altos City ordinance:

- prohibits installation of small cells on public utility easements in residential neighborhoods
- 500 foot setbacks for small cells for multi-family residences in commercial districts
- 500 ft separation from schools
- 1500 ft separation between nodes

San Diego County, California

- "SCWs shall not be located within 1,000 feet of schools, child care centers, hospitals, or churches."

CONNECTICUT

- Easton CN City Council passed a 5G cease and desist resolution
- Warren, Connecticut Policy defines "adequate coverage" and "adequate capacity." and was designed "to locate towers and/or antennas in a manner which protects property values, as well as the general safety, health, welfare and quality of life of the citizens." Coverage is considered to be "adequate" within that area surrounding a Base Station where the predicted or measured median field strength of the transmitted signal is such that the majority of the time, transceivers properly installed and operated will be able to communicate with the base station.

FLORIDA

- Coconut Creek FL Commission adopted a Resolution on 5G and radiofrequency radiation.
- Hallandale Beach FL Resolution urges the federal government to initiate independent health studies on 5G.
- Lavallette FL Resolution 2021-58: Applicant shall obtain certification from the Federal Aviation Administration and the United States Dept. of Defense demonstrating that the installation does not emit RF frequencies which may interfere with avionics of any approaching civil or military aircraft." The City also requires the applicant to provide RF meters used by their technicians and train City employees. Verizon cannot install more than a total of 20 "small cell" nodes throughout the Borough to support 5G.

HAWAII

- Hawai'i County Council passed a Resolution to halt 5G

ILLINOIS

- Oak Brook IL Resolution calls for local control re small cells.

INDIANA

Carmel City IN Council resolution asks state lawmakers, FCC and Congress to limit 5G until health effects fully understood.

MASSACHUSETTS

Randolph MA 500 ft setback. Yearly RFR measurements. Lunenburg and Great Barrington MA 500 ft setback Stockbridge MA prohibits a tower from being built 1000 feet from a school, park or athletic field and 600 ft from residence.

NEW JERSEY

- Little Silver, NJ Carriers should provide notice to property owners within 500 feet of proposed facility.

NEW YORK

- Scarsdale NY: 500 foot setbacks to homes preferred.
- Copake NY: Pre/post testing by RF engineer. No repeater closer than 200 ft to dwelling. No tower closer than 1500 ft to residence/church.
- Community Boards issuing Moratoriums on 5G poles

NEW HAMPSHIRE

- Proposed State Bill - 1640 ft setbacks.
- Keene NH Resolution to halt 5G
- Bedford NH 750 ft. setback

OHIO

- Mason OH Zoning Ordinance No small cells in residential areas or within 100 feet of residential prop; 2000 feet apart (unless colocated); equipment should be underground or wholly contained.

OKLAHOMA

- Sallisaw OK 1,500 feet setback

TENNESSEE

- Farragut City Resolution to halt 5G

WISCONSIN

- Greendale WI passed Resolution R2018-20 referring to the FCC's actions stripping local authority as "an unprecedented attack on local control."

5G & CELL TOWERS



EUROPE

- Resolutions to halt 5G in numerous European cities including Trafford, UK, Lille, France, Ormidia, Cyprus, Councils in Ireland and more.

ITALY

- 600 municipalities have passed resolution to halt 5G.

UNITED STATES

- Los Angeles CA Public Schools: RFR Limit 10,000x less than FCC.
- Resolutions to halt 5G passed in Hawaii County HI, Farragut TN, Keene NH & Easton CT.
- Numerous cities restrict cell antennas near homes including: Los Altos, Petaluma, Mill Valley, Malibu and San Diego County CA, Bedford NH and more.
- New Hampshire 5G Commission's 15 Recommendations include increasing transparency, reduce public exposure, research health effects and protect wildlife and trees.
- Oregon investigating health effects of wireless.
- Palo Alto, Los Angeles LA Schools Greenbelt MD, Bar Harbor ME; No school cell towers

CHILE

- Cell antennas prohibited in "sensitive areas" - kindergartens, hospitals and nursing homes.

BANGLADESH

- No cell towers on homes, schools, colleges, playing fields, populated areas and heritage areas.

FRANCE

- 60 mayors/officials petition to halt 5G.
- Federal health agency investigating 5G
- 5G antenna RFR is measured.

SWITZERLAND

- Parliament refused to weaken radio frequency radiation (RFR) limits after 5G Report.

NETHERLANDS

- Health Council recommends against 26 GHz for 5G due to lack of safety data.

RUSSIA

- No cell towers near schools.

ISRAEL

- Cell tower setback 100m from schools/ homes.

CANADA

- City of Toronto "Prudent Avoidance Policy" for Cell Towers.

BULGARIA

- Mezdra and Balchik have banned 5G.

GREECE

- The installation of cell towers at the premises of schools, kindergartens, hospitals or eldercare facilities is prohibited.

CYPRUS

- Cyprus National Committee on Environment and Child Health 5G Position Paper calls for 5G free zones.

AUSTRALIA


- New South Wales Dept. of Education policy objects to towers on/near schools.

LITHUANIA

- Cell antennas prohibited on kindergartens and hospitals.

INDIA

- RFR limit tightened to 1/10 of CNIRP limits after Inter-Ministerial Report on impacts to wildlife.
- Mumbai, Zilla Parishad & Karnataka: Cell towers prohibited/removed near schools, colleges, orphanages and old age homes.
- Brihanmumbai Municipal: Cell towers banned at parks and playgrounds.
- State of Rajasthan: Supreme Court of India upheld removal of "hazardous to life" cell towers from vicinity of schools, hospitals/playgrounds.



WIRELESS RADIATION AND CHILDREN'S HEALTH

EXPERT VOICES



"The National Toxicology Program studies clearly showed that non-ionizing cell phone radiofrequency radiation can cause cancers and other adverse health effects. An important lesson that should be learned is that we cannot assume any current or future wireless technology such as 5G is safe without adequate testing."

— **Ronald Melnick PhD 28 year scientist at National Institutes of Health**

"I recommend public health organizations raise awareness and educate the public on why and how to reduce our daily exposure to wireless radio frequency radiation. Protective public health policy is needed now. It is time for regulatory bodies to fully evaluate the research and develop science based exposure limits that truly protect the public and the environment."

— **Linda S. Birnbaum, PhD, Former Director, National Institute of Environmental Health Sciences and National Toxicology Program of the National Institutes of Health.**

"Given the human, animal and experimental evidence, I assert that, to a reasonable degree of scientific certainty, the probability that RF exposure causes gliomas and neuromas is high."

— **Christopher Portier PhD former Director of the United States National Center for Environmental Health at the CDC, former Director of the U.S. Agency for Toxic Substances and Disease Registry.**

I am calling on my industry to bring safer technology to market. The current implementation of technology is not safe. Take a good look at the science. This is about our children's future. Do not be lulled into believing that 25-year-old standards can protect the youngest and most vulnerable. They simply cannot."

— **Frank Clegg, Former President of Microsoft Canada, CEO of Canadians for Safe Technology**

"Most parents believe that cellphones were safety-tested before they came on the market. We assume that our federal health and environmental agencies regularly review the latest research and ensure that these incredible devices are safe. They do not. As we sadly learned with early childhood lead exposures leaving long-lasting impairments, the developing brain is particularly susceptible."

— **Jerome Paulson, MD , Professor Emeritus, George Washington University, Milliken School of Public Health, former Chair of American Academy of Pediatrics Committee on Environmental Health**

PARENT TEACHER ASSOCIATIONS

OPPOSE CELL TOWERS

Agenda No. 11A Attachment B
Page 46 of 186
Exhibit A

August 6, 2018

Pacific Grove City Council
City Hall
300 Forest Ave.
Pacific Grove, CA 93950

Dear members of Pacific Grove City Council,

I am writing you on behalf of Pacific Grove High School PTA in regard to the Pacific Grove Planning Commission's vote on July 26, 2018, which approved a request by Verizon Wireless to install and maintain a cell tower adjacent to Pacific Grove High School (PGHS). For the reasons described below, the Pacific Grove High School PTA **is strongly opposed** to the location of the Verizon cell tower and is requesting that the City Council consider and support the appeal that is being filed by a group of concerned parents who live in Pacific Grove and send their children to PG schools.

The installment of a cell tower adjacent to PGHS poses significant potential health dangers to both students and staff at PGHS. While some argue that radiation emitted from a cell tower is not a health danger, data from many studies indicate the opposite. Research shows that children and pregnant women are the most vulnerable – two demographics most likely to be on PGHS school grounds on a regular basis. The actual placement of the cell tower – near the back of PGHS and very close to Forest Grove Elementary School – only increases the concerns of the frequency in exposure.

Cell towers also pose a risk to students due to fire hazard. Many cell towers throughout the United States have caught fire and collapsed, posing a significant safety concern, especially in an area with young students walking to and from school every day.

The mission of all PTAs nationwide is to make every child's potential a reality by engaging and empowering families and communities to advocate for all children. Our local PTA is very active in expressing our support for or opposition to issues dealing with the health, safety, education, or general well-being of children and youth in our community.

The members of the Pacific Grove High School PTA strongly urge you to please reconsider the Pacific Grove Planning Commission's previous vote and rescind approval for the Verizon cell tower at Pacific Grove High School.

Sincerely,

Julie Kavanaugh
President, Pacific Grove High School PTA

Health impact of 5G

Current state of knowledge of 5G-related carcinogenic and reproductive/developmental hazards as they emerge from epidemiological studies and *in vivo* experimental studies

The upcoming deployment of 5G mobile networks will allow for significantly faster mobile broadband speeds and increasingly extensive mobile data usage. Technical innovations include a different transmission system (MIMO: use of multiple-input and multiple-output antennas), directional signal transmission or reception (beamforming), and the use of other frequency ranges. At the same time, a change is expected in the exposure to electromagnetic fields (EMF) of humans and the environment. In addition to those used to date, the 5G pioneer bands identified at EU level have frequencies of 700 MHz, 3.6 GHz (3.4 to 3.8 GHz) and 26 GHz (24.25 to 27.5 GHz). The first two frequencies (FR1) are similar to those used for 2G to 4G technologies and have been investigated in both epidemiological and experimental studies for different end points (including carcinogenicity and reproductive/developmental effects), while 26 GHz (FR2) and higher frequencies have not been adequately studied for the same end points.

The International Agency for Research on Cancer (IARC) classified radiofrequency (RF) EMF as 'possibly carcinogenic to humans' (Group 2B) and recently recommended RF exposure for re-evaluation 'with high priority' (IARC, 2019). Since 2011 a great number of studies have been performed, both epidemiological and experimental. The present review addresses the current knowledge regarding both carcinogenic and reproductive/developmental hazards of RF as exploited by 5G. There are various *in vivo* experimental and epidemiological studies on RF at a lower frequency range (450 to 6000 MHz), which also includes the frequencies used in previous generations' broadband cellular networks, but very few (and inadequate) on the higher frequency range (24 to 100 GHz, centimetre/MMW).

The review shows: 1) 5G lower frequencies (700 and 3 600 MHz): a) limited evidence of carcinogenicity in epidemiological studies; b) sufficient evidence of carcinogenicity in experimental bioassays; c) sufficient evidence of reproductive/developmental adverse effects in humans; d) sufficient evidence of reproductive/developmental adverse effects in experimental animals; 2) 5G higher frequencies (24.25-27.5 GHz): the systematic review found no adequate studies either in humans or in experimental animals.

Conclusions: 1) cancer: FR1 (450 to 6 000 MHz): EMF are probably carcinogenic for humans, in particular related to gliomas and acoustic neuromas; FR2 (24 to 100 GHz): no adequate studies were performed on the higher frequencies; 2) reproductive developmental effects: FR1 (450 to 6 000 MHz): these frequencies clearly affect male fertility and possibly female fertility too. They may have possible adverse effects on the development of embryos, foetuses and newborns; FR2 (24 to 100 GHz): no adequate studies were performed on non-thermal effects of the higher frequencies.

SCIENTIFIC RESEARCH STUDIES



European Parliament requested a research report "[Health Impact of 5G](#)" which was released in July 2021 and concluded that commonly used RFR frequencies (450 to 6000 MHz) are probably carcinogenic for humans and clearly affect male fertility with possible adverse effects on the development of embryos, fetuses and newborns.

A review entitled "[Evidence for a health risk by RF on humans living around mobile phone base stations: From radiofrequency sickness to cancer](#)" reviewed the existing scientific literature and found radiofrequency sickness, cancer and changes in biochemical parameters ([Balmori 2022](#)).

A [study](#) published in Electromagnetic Biology and Medicine found changes in blood considered biomarkers predictive of cancer in people living closer to cell antenna arrays ([Zothansiana 2017](#)).

A [study](#) published in the International Journal of Environmental Research and Public Health found higher exposure to cell network arrays linked to higher mortality from all cancer and specifically lung and breast cancer ([Rodrigues 2021](#)).

A 10-year [study](#) published in Science of the Total Environment on cell phone network antennas by the local Municipal Health Department and several universities in Brazil found a clearly elevated relative risk of cancer mortality at residential distances of 500 meters or less from cell phone towers ([Dode 2011](#)).

A [study](#) commissioned by the Government of Styria, Austria found a significant cancer incidence in the area around the RF transmitter as well as significant exposure-effect relationships between radiofrequency radiation exposure and the incidence of breast cancers and brain tumors ([Oberfeld 2008](#)).

A [review](#) published in Experimental Oncology found "alarming epidemiological and experimental data on possible carcinogenic effects of long term exposure to low intensity microwave (MW) radiation." A year of operation of a powerful base transmitting station for mobile communication reportedly resulted in a dramatic increase of cancer incidence among the population living nearby ([Yakymenko 2011](#)).



Bold blue on this PDF are hyperlinked.

CHILDREN'S VULNERABILITY TO WIRELESS RADIOFREQUENCY (RF) RADIATION



The American Academy of Pediatrics states:

"In recent years, concern has increased about exposure to radio frequency (RF) electromagnetic radiation emitted from cell phones and phone station antennas. An Egyptian study confirmed concerns that living nearby mobile phone base stations increased the risk for developing:

- Headaches
- Memory problems
- Dizziness
- Depression
- Sleep problems

Short-term exposure to these fields in experimental studies have not always shown negative effects, but this does not rule out cumulative damage from these fields, so larger studies over longer periods are needed to help understand who is at risk. In large studies, an association has been observed between symptoms and exposure to these fields in the everyday environment."

-American Academy of Pediatrics
[HealthyChildren.org](https://www.healthychildren.org)

Cell towers and cell phones emit wireless radiofrequency (RF) radiation.

Children are more vulnerable to RF radiation, just as they are to other environmental exposures. They have proportionately more exposures to RF compared to adults. More importantly, even very low exposures to children can have serious impacts later in life because their nervous and immune systems are still in development.

Children absorb higher levels of RF radiation deeper into their brains and bodies because they have:

- Thinner skulls allow RF radiation to move easier into the brain.
- Higher water content in brain tissue which is more conductive to electricity.
- Smaller heads result in a shorter distance for the RF to travel from the skull to critical brain regions important for learning and memory.

Children are more sensitive to RF impacts because:

- Their brains are still developing.
- Children have more active stem cells- a type of cell scientifically found to be uniquely impacted by RF.
- Children will have a longer lifetime of higher exposures, starting from before they are born.

Bold blue on this PDF are hyperlinked.

ENVIRONMENTAL HEALTH TRUST | [EHTRUST.ORG](https://ehtrust.org)

CELL TOWER RF RADIATION AND CANCER

International Agency for Research on Cancer



World Health
Organization

PRESS RELEASE
N° 208

31 May 2011

IARC CLASSIFIES RADIOFREQUENCY ELECTROMAGNETIC FIELDS AS POSSIBLY CARCINOGENIC TO HUMANS

Lyon, France, May 31, 2011 -- The WHO/International Agency for Research on Cancer (IARC) has classified radiofrequency electromagnetic fields as possibly carcinogenic to humans (Group 2B), based on an increased risk for glioma, a malignant type of brain cancer¹, associated with wireless phone use.

The World Health Organization International Agency for Research on Cancer Classified Radiofrequency Radiation as a "Possible" Carcinogen in 2011

In 2011, radiofrequency electromagnetic fields (RF-EMF) were classified as a Group 2B possible carcinogen by the World Health Organization's International Agency for Research on Cancer (WHO/IARC).

The WHO/IARC scientists clarified that this determination was for RF-EMF from any source be it cell phones, wireless devices, cell towers or any other type of wireless equipment.


Since 2011, the published peer-reviewed scientific evidence associating RF-EMF (also known as RF-EMR and RFR) to cancer and other adverse effects has significantly increased.

A large-scale animal study published in Environmental Research found rats exposed to RF levels comparable to cell tower emissions had elevated cancers, the very same cancers also found in the US National Toxicology Program animal study of cell phone level RF that found "clear evidence" of cancer in carefully controlled conditions (Falcioni 2018).

In 2019, the WHO/IARC advisory committee recommended that radiofrequency radiation be re-evaluated as a "high" priority in light of the new research. The date of the re-evaluation has not been set.

Currently, several scientists conclude that the weight of currently available, peer-reviewed evidence supports the conclusion that radiofrequency radiation is a proven human carcinogen (Hardell and Carlberg 2017, Peleg et al. 2022, Miller et al. 2018).

PUBLISHED RESEARCH STUDIES



RESEARCHERS RECOMMEND CELL TOWERS BE DISTANCED AWAY FROM HOMES AND SCHOOLS

The review paper entitled “[Limiting liability with positioning to minimize negative health effects of cellular phone towers](#)” reviewed the “large and growing body of evidence that human exposure to RFR from cellular phone base stations causes negative health effects.” The authors recommend restricting antennas near homes, and restricting antennas within 500 meters of schools and hospitals to protect companies from future liability ([Pearce 2020](#)).

An [analysis](#) of 100 studies published in *Environmental Reviews* found approximately 80% showed biological effects near towers. “As a general guideline, cell base stations should not be located less than 1500 ft from the population, and at a height of about 150 ft” ([Levitt 2010](#)).

A [review](#) published in the *International Journal of Occupational and Environmental Health* found people living less than 500 meters from base station antennas had increased adverse neuro-behavioral symptoms and cancer in eight of the ten epidemiological studies ([Khurana 2010](#)).

A [paper](#) by human rights experts published in *Environment Science and Policy* documented the accumulating science indicating safety is not assured, and considered the issue within a human rights framework to protect vulnerable populations from environmental pollution. “We conclude that, because scientific knowledge is incomplete, a precautionary approach is better suited to State obligations under international human rights law” ([Roda and Perry 2014, PDF](#)).

APARTMENTS & CONDO BUILDINGS

INCREASED RF RADIATION FROM CELL ANTENNAS



The study "[Radiofrequency radiation from nearby mobile phone base stations-a case comparison of one low and one high exposure apartment](#)" published in *Oncology Letters* by [Koppel et al. \(2019\)](#) measured 2 apartments and found that the apartment with high RF levels had outdoor areas as close as 6 meters (about 19.6 feet) from transmitting base station cell antennas. In contrast, the apartment with low RF exposure had cell antennas at 40 meters (about 131 feet) away from the balcony.

Furthermore, the researchers also found that both high- and low-RF apartments had good mobile phone reception, and they concluded, "therefore, installation of base stations to risky places cannot be justified using the good reception requirement argument."

A measurement study by [Baltrėnas et al. \(2012\)](#) published in *Journal of Environmental Engineering and Landscape Management* investigated RF power density levels from cell phone antennas located 35 meters away from a 10-story apartment building. The transmitting antennas were approximately at the same height as the 6th floor of the building. The researchers found the highest RF levels at floors 5, 6 and 7. The RF at the 6th floor balcony was three times higher than the 3rd floor balcony. The RF power density at the 6th floor was about 15 times the RF measured at the first floor.

A [case report by Hardell et al. \(2017\)](#) of RF levels in an apartment in close proximity to rooftop cellular network antennas used an exposimeter to measure levels of different types of RF in the apartment and balconies including TV, FM, TETRA emergency services, 2G GSM, 3G UMTS, 4G LTE, DECT cordless, Wi-Fi 2.4 GHz and 5 GHz and WiMAX. The closest transmitting antennas were 6 meters away from the balcony. The researchers found 97.9% of the mean RF radiation was caused by downlink from the 2G, 3G and 4G base stations. (Downlink means frequencies emitted "down" from the base station cellular antennas.) The researchers found that if the base station RF emissions were excluded, the RF radiation in the children's bedrooms was reduced approximately 99%.

The researchers conclude, "due to the current high RF radiation, the apartment is not suitable for long-term living, particularly for children who may be more sensitive than adults."

INCREASED EXPOSURE FROM 5G/4G "SMALL" CELL ANTENNAS LOCATED CLOSE TO PEOPLE

A study entitled "[Very high radiofrequency radiation at Skeppsbron in Stockholm, Sweden from mobile phone base station antennas positioned close to pedestrians' heads](#)" published in *Environmental Research* by Koppel et al. (2022) created an RF heat map of RF measurements, finding that the highest RF measurements were in areas of close proximity to the base station antennas. The researchers concluded with recommendations to reduce close proximity placements such as positioning antennas "as far as possible from the general public" like in high-elevation locations or more remote areas.

A study entitled "[Measurements of radiofrequency electromagnetic fields, including 5G, in the city of Columbia, South Carolina, USA](#)" published in the *World Academy of Sciences Journal* found the highest RF levels in areas where the cell phone base station antennas were placed on top of utility poles, street lamps, traffic lights or other posts near to the street. The scientists compared their [2022 findings](#) to an earlier [2019 published review](#) on the mean outdoor exposure level of European cities and they found the South Carolina measurements to be higher.

The researchers concluded that the highest exposure areas were due to two reasons: cell phone base antennas on top of high-rise buildings provide "good cell coverage reaching far away, but creating elevated exposure to the radiofrequency electromagnetic fields at the immediate vicinity; and cell phone base station antennas installed on top of utility poles have placed the radiation source closer to humans walking on street level."



Figure 7. Gervais Street: Cell phone base station antenna placed close to street level and causing high exposure to pedestrians and nearby café visitors (exposure scenario illustration). The antenna appears camouflaged and seemingly part of a utility pole. The measurer only discovered the antenna due to the high radiofrequency levels in the vicinity.

HEALTH SYMPTOMS REPORTED BY PEOPLE LIVING CLOSE TO CELL ANTENNAS

Image: Figure 1: Top floor apartment adjacent to base stations. Nilsson M, Hardell L. (2023) Development of the Microwave Syndrome in Two Men Shortly after Installation of 5G on the Roof above their Office. Ann Clin Case Rep



RESEARCH ON ANTENNAS CLOSE TO HOMES, SCHOOL AND WORK

Surveys of people living near cell tower antennas in [France](#), [Spain](#), [Iraq](#), [India](#), [Germany](#), [Egypt](#), [Poland](#) have found significantly higher reports of health issues including sleep issues, fatigue and headaches (See [Santini et al. 2003](#), [López 2021](#), [Alazawi 2011](#), [Pachua and Pachua 2016](#), [Eger et al. 2004](#), [Abdel-Rassoul et al. 2007](#), [Bortkiewicz et al., 2004](#)).

A [study](#) published in *American Journal of Men's Health* linked higher cell tower RFR exposures to delayed fine and gross motor skills and to deficits in spatial working memory and attention in school adolescents ([Meo 2018](#)).

A [study](#) published in *Environmental Research and Public Health* found higher exposures linked to higher risk of type 2 diabetes ([Meo 2015](#)).

A study following people for 6 years linked increased cell phone and cell phone tower antenna exposure to altered levels of hormones including cortisol, thyroid, prolactin and testosterone ([Eskander et al. 2021](#)).

A [study](#) that followed people in a German town after a cell tower was erected found stress hormones adrenaline and noradrenaline significantly increased over the first 6 months after the antenna activation and decreased dopamine and PEA levels after 18 months ([Buchner 2011](#)).

Two published case report document illness that developed after 5G antennas were installed. In [Hardell and Nilsson 2023](#), a couple developed microwave syndrome symptoms (e.g., neurological symptoms, tinnitus, fatigue, insomnia, emotional distress, skin disorders, and blood pressure variability) after a 5G base station was installed on the roof above their apartment.

Similarly, in "[Development of the Microwave Syndrome in Two Men Shortly after Installation of 5G on the Roof above their Office](#)" two men developed symptoms after 5G antennas were activated on the roof of their workplace. The symptoms disappeared in both men within a couple of weeks (case 1) or immediately (case 2) after leaving the office.

PUBLISHED RESEARCH ON 5G



New York City Jumbo 5G poles with 5 tiers to house transmitting antennas from numerous carriers.



New York City "small cell" antennas in front of living room window.

Scientists state that 5G's higher frequencies cannot be assumed safe.

5G systems are using low band frequencies well associated with harmful effects ([ICBE-EMF 2022](#), [European Parliament 2021](#), [Panagopoulos et al. 2021](#)). However 5G networks are also using higher frequencies such as 3.5 GHz and into the mmWave range with 24 GHz and higher.

Contrary to claims that the 5G's higher frequencies simply "bounce" off the skin, researchers have documented that the coiled portion of the skin's sweat duct can be regarded as a helical antenna in the sub-THz band and the skin, our largest organ, can intensely absorb the higher 5G frequencies ([Feldman and Ben Ishai 2017](#)).

Reviews of 5G health effects caution that the expected real-world impact would be far more serious due to the complex waveforms and other combinations with other toxic stimuli in the environment ([Kostoff et al 2020](#), [Russell, 2018](#), [Belyaev 2019](#), [McCredden et al 2023](#)).

Researchers will often experiment with zebrafish, rodents and fruit flies to gain data on potential health effects to humans. An Oregon State University study on zebrafish exposed to 3.5 GHz ([Dasgupta et al. 2022](#)) found "significant abnormal responses in RFR-exposed fish" which "suggest potential long-term behavioral effects. Yang et al 2022 found 3.5 GHz induced oxidative stress in guinea pigs.

A study on 3.5 GHz exposure to both diabetic and healthy rats ([Bektas et al 2022](#)) found an increase in degenerated neurons in the hippocampus of the brains, changes in oxidative stress parameters and changes in the energy metabolism and appetite of both healthy and diabetic rats. The researchers conclude that, "5G may not be innocent in terms of its biological effects, especially in the presence of diabetes."

REFERENCES/CITATIONS

- Abdel-Rassoul, G., et al (2007). [Neurobehavioral effects among inhabitants around mobile phone base stations](#). *NeuroToxicology*
- Balmori A. (2002) [Evidence for a health risk by RF on humans living around mobile phone base stations: From radiofrequency sickness to cancer](#). *Environmental Research*
- Dode, A. C et al (2011). [Mortality by neoplasia and cellular telephone base stations in the Belo Horizonte municipality, Minas Gerais state, Brazil](#). *Science of The Total Environment*
- Hardell, L., & Carlberg, M. (2019). [Comments on the US National Toxicology Program technical reports on toxicology and carcinogenesis study in rats exposed to whole-body radiofrequency radiation at 900 MHz and in mice exposed to whole-body radiofrequency radiation at 1,900 MHz](#). *International Journal of Oncology*
- Hardell, L., & Koppel, T. (2022). [Electromagnetic hypersensitivity close to mobile phone base stations – a case study in Stockholm, Sweden](#). *Reviews on Environmental Health*.
- Khurana et al. (2010). [Epidemiological evidence for a health risk from mobile phone base stations](#). *International Journal of Occupational and Environmental Health*
- Koppel et al (2022). [Very high radiofrequency radiation at Skeppsbron in Stockholm, Sweden from mobile phone base station antennas positioned close to pedestrians' heads](#). *Environmental Research*
- Levitt & Lai, H. (2011). [Corrigendum: Biological effects from exposure to electromagnetic radiation emitted by cell tower base stations and other antenna arrays](#). *Environmental Reviews*
- López et al (2021). [What is the radiation before 5G? A correlation study between measurements in situ and in real time and epidemiological indicators in Vallecas, Madrid](#). *Environmental Research*
- Meo et al (2019). [Mobile Phone Base Station Tower Settings Adjacent to School Buildings: Impact on Students' Cognitive Health](#). *American Journal of Men's Health*
- Meo et al (2015a). [Association of Exposure to Radio-Frequency Electromagnetic Field Radiation \(RF-EMFR\) Generated by Mobile Phone Base Stations with Glycated Hemoglobin \(HbA1c\) and Risk of Type 2 Diabetes Mellitus](#). *International Journal of Environmental Research and Public Health*
- Pearce, J. M. (2020). [Limiting liability with positioning to minimize negative health effects of cellular phone towers](#). *Environmental Research*
- Roda, C., & Perry, S. (2014). [Mobile phone infrastructure regulation in Europe: Scientific challenges and human rights protection](#). *Environmental Science & Policy*
- Rodrigues et al (2021). [The Effect of Continuous Low-Intensity Exposure to Electromagnetic Fields from Radio Base Stations to Cancer Mortality in Brazil](#). *International Journal of Environmental Research and Public Health*
- Santini et al. (2003). [Survey Study of People Living in the Vicinity of Cellular Phone Base Stations](#). *Electromagnetic Biology and Medicine*
- Thamilselvan et al (2021) [Micronuclei analysis in people residing within 25 m of radiation-exposed areas around mobile towers in Chennai, India: An observational study](#). *Journal of International Oral Health*
- Yakymenko et al (2011). [Long-term exposure to microwave radiation provokes cancer growth: Evidences from radars and mobile communication systems](#). *Experimental Oncology*
- Zothansiamia et al (2017). [Impact of radiofrequency radiation on DNA damage and antioxidants in peripheral blood lymphocytes of humans residing in the vicinity of mobile phone base stations](#). *Electromagnetic Biology and Medicine*
- Belyaev et al (2022) [Possible health risks from exposure to microwaves from base stations](#), Conference Paper Department of Radiobiology, Cancer Research Institute, Biomedical Research Center *Conference paper

REFERENCES/CITATIONS

5G

- Betzalel et al. (2018). [The human skin as a sub-THz receiver—Does 5G pose a danger to it or not?](#) *Environmental Research*
- Betzalel et al. (2017). [The Modeling of the Absorbance of Sub-THz Radiation by Human Skin](#). *IEEE Transactions on Terahertz Science and Technology*
- Dasgupta et al. (2022). [Transcriptomic and Long-Term Behavioral Deficits Associated with Developmental 3.5 GHz Radiofrequency Radiation Exposures in Zebrafish](#). *Environmental Science & Technology Letters*
- Di Ciaula, A. (2018). [Towards 5G communication systems: Are there health implications?](#) *International Journal of Hygiene and Environmental Health*
- Frank, J. W. (2021). [Electromagnetic fields, 5G and health: What about the precautionary principle?](#) *Journal of Epidemiology and Community Health*
- Hardell, L., & Carlberg, M. (2020). [\[Comment\] Health risks from radiofrequency radiation, including 5G, should be assessed by experts with no conflicts of interest](#). *Oncology Letters*
- Hardell, L., & Nilsson, M. (2023). [Case Report: The Microwave Syndrome after Installation of 5G Emphasizes the Need for Protection from Radiofrequency Radiation](#). *Annals of Case Reports*.
- Nilsson M, Hardell L. (2023) [Development of the Microwave Syndrome in Two Men Shortly after Installation of 5G on the Roof above their Office](#). *Ann Clin Case Rep*. 8: 2378.
- Hinrikus et al. (2022). [Possible health effects on the human brain by various generations of mobile telecommunication: A review based estimation of 5G impact](#). *International Journal of Radiation Biology*
- Kostoff et al. (2020). [Adverse health effects of 5G mobile networking technology under real-life conditions](#). *Toxicology Letters*
- Nasim, I., & Kim, S. (2019). [Adverse Impacts of 5G Downlinks on Human Body](#). *2019 SoutheastCon IEEE*
- Russell, C. L. (2018). [5G wireless telecommunications expansion: Public health and environmental implications](#). *Environmental Research*
- Yang et al. (2022). [Effects of Acute Exposure to 3500 MHz \(5G\) Radiofrequency Electromagnetic Radiation on Anxiety-Like Behavior and the Auditory Cortex in Guinea Pigs](#). *Bioelectromagnetics*

Increasing Exposures From Expanding 5G Networks and Close Proximity “Small Cell” Antennas

- Baltrėnas et al.(2012). [Research and evaluation of the intensity parameters of electromagnetic fields produced by mobile communication antennas](#). *Journal of Environmental Engineering and Landscape Management*
- Bhatt et al. (2017). [Radiofrequency-electromagnetic field exposures in kindergarten children](#). *Journal of Exposure Science & Environmental Epidemiology*
- Bonato et al. (2022). [Computational Assessment of RF Exposure Levels due to 5G Mobile Phones](#). *2022 Microwave Mediterranean Symposium*
- Carlberg et al. (2019). [High ambient radiofrequency radiation in Stockholm city, Sweden](#). *Oncology Letters*
- El-Hajj et al. (2020). [Radiation Analysis in a Gradual 5G Network Deployment Strategy](#). *2020 IEEE 3rd 5G World Forum (5GWF)*
- Hardell et al. (2018). [Radiofrequency radiation from nearby base stations gives high levels in an apartment in Stockholm, Sweden: A case report](#). *Oncology Letters*
- Hardell et al. (2017). [High radiofrequency radiation at Stockholm Old Town: An exposimeter study including the Royal Castle, Supreme Court, three major squares and the Swedish Parliament](#). *Molecular and Clinical Oncology*
- Hardell, L., Koppel, T., Carlberg, M., Ahonen, M., & Hedendahl, L. (2016). [Radiofrequency radiation at Stockholm Central Railway Station in Sweden and some medical aspects on public exposure to RF fields](#). *International Journal of Oncology*
- Koppel et al. (2022). [Very high radiofrequency radiation at Skeppsbron in Stockholm, Sweden from mobile phone base station antennas positioned close to pedestrians' heads](#). *Environmental Research*
- Koppel et al. (2019). [Radiofrequency radiation from nearby mobile phone base stations—a case comparison of one low and one high exposure apartment](#). *Oncology Letters*

REFERENCES/CITATIONS

Koppel, T., & Hardell, L. (2022). [Measurements of radiofrequency electromagnetic fields, including 5G, in the city of Columbia, SC, USA](#). *World Academy of Sciences Journal*

Mazloum et al. (2019). [RF-EMF exposure induced by mobile phones operating in LTE small cells in two different urban cities](#). *Annals of Telecommunications*

Urbinello et al. (2014). [Temporal trends of radio-frequency electromagnetic field \(RF-EMF\) exposure in everyday environments across European cities](#). *Environmental Research*, 134, 134–142.

4G LTE

Broom et al. (2019). [Early-Life Exposure to Pulsed LTE Radiofrequency Fields Causes Persistent Changes in Activity and Behavior in C57BL/6 J Mice](#). *Bioelectromagnetics*

Choi et al. (2020). [Continuous Exposure to 1.7 GHz LTE Electromagnetic Fields Increases Intracellular Reactive Oxygen Species to Decrease Human Cell Proliferation and Induce Senescence](#). *Scientific Reports*

Lv et al. (2014). [The alteration of spontaneous low frequency oscillations caused by acute electromagnetic fields exposure](#). *Clinical Neurophysiology*

Malik et al. (2021). [Short- and long-duration exposures to cell-phone radiofrequency waves produce dichotomous effects on phototactic response and circadian characteristics of locomotor activity rhythm in zebrafish, Danio rerio](#). *Biological Rhythm Research*

Oh, J. J., Byun, S.-S., Lee, S. E., Choe, G., & Hong, S. K. (2018). [Effect of Electromagnetic Waves from Mobile Phones on Spermatogenesis in the Era of 4G-LTE](#). *BioMed Research International*, 2018, 1801798.

Özdemir et al. (2021). [The effect of 4.5 G \(LTE Advanced-Pro network\) mobile phone radiation on the optic nerve](#). *Cutaneous and Ocular Toxicology*

Souffi et al. (2022). [Exposure to 1800 MHz LTE electromagnetic fields under proinflammatory conditions decreases the response strength and increases the acoustic threshold of auditory cortical neurons](#). *Scientific Reports*

Wei et al. (2019). [Modulation of resting-state brain functional connectivity by exposure to acute fourth-generation long-term evolution electromagnetic field: An fMRI study](#). *Bioelectromagnetics*

Yang et al. (2021). [Functional and network analyses of human exposure to long-term evolution signal](#). *Environmental Science and Pollution Research International*

Yang et al. (2017). [Long-Term Evolution Electromagnetic Fields Exposure Modulates the Resting State EEG on Alpha and Beta Bands](#). *Clinical EEG and Neuroscience*

Yu et al. (2020). [Long-term exposure to 4G smartphone radiofrequency electromagnetic radiation diminished male reproductive potential by directly disrupting Spock3–MMP2–BTB axis in the testes of adult rats](#). *Science of The Total Environment*

WILDLIFE , WIRELESS AND NON IONIZING ELECTROMAGNETIC FIELDS

SCIENTIFIC RESEARCH

REVIEWS

Balmori, A. (2015). Anthropogenic radiofrequency electromagnetic fields as an emerging threat to wildlife orientation. *Science of The Total Environment*, 518–519, 58–60.

Balmori A. (2014). Electrosmog and species conservation. *Science of The Total Environment*, 496:314-316

Cucurachi, S., Tamis, W. L. M., Vijver, M. G., Peijnenburg, W. J. G. M., Bolte, J. F. B., & de Snoo, G. R. (2013). A review of the ecological effects of radiofrequency electromagnetic fields (RF-EMF). *Environment International*, 51, 116–140.

Levitt BB, Lai HC and Manville AM II (2022) Low-level EMF effects on wildlife and plants: What research tells us about an ecosystem approach. *Front. Public Health* 10:1000840. doi: 10.3389/fpubh.2022.1000840

Levitt, B. B., Lai, H. C., & Manville, A. M. (2021). Effects of non-ionizing electromagnetic fields on flora and fauna, Part 3. Exposure standards, public policy, laws, and future directions. *Reviews on Environmental Health*.

Levitt, B. B., Lai, H. C., & Manville, A. M. (2022a). Effects of non-ionizing electromagnetic fields on flora and fauna, part 1. Rising ambient EMF levels in the environment. *Reviews on Environmental Health*, 37(1), 81–122.

Levitt, B. B., Lai, H. C., & Manville, A. M. (2022b). Effects of non-ionizing electromagnetic fields on flora and fauna, Part 2 impacts: How species interact with natural and man-made EMF. *Reviews on Environmental Health*, 37(3), 327–406.

BIRDS

Balmori A. (2022). Corneal opacity in Northern Bald Ibises (*Geronticus eremita*) equipped with radio transmitters. *Electromagnetic Biol Med.*174-176.

Fernie, K. J., & Reynolds, S. J. (2005). The effects of electromagnetic fields from power lines on avian reproductive biology and physiology: A review. *Journal of Toxicology and Environmental Health. Part B, Critical Reviews*, 8(2), 127–140

Schwarze, S., Schneider, N.-L., Reichl, T., Dreyer, D., Lefeldt, N., Engels, S., Baker, N., Hore, P. J., & Mouritsen, H. (2016). Weak Broadband Electromagnetic Fields are More Disruptive to Magnetic Compass Orientation in a Night-Migratory Songbird (*Erithacus rubecula*) than Strong Narrow-Band Fields. *Frontiers in Behavioral Neuroscience*, 10.

Tonelli, B. A., Youngflesh, C., & Tingley, M. W. (2023). Geomagnetic disturbance associated with increased vagrancy in migratory landbirds. *Scientific Reports*, 13(1), Article 1.

Wiltshko, R., Thalau, P., Gehring, D., Nießner, C., Ritz, T., & Wiltshko, W. (2015). Magnetoreception in birds: The effect of radio-frequency fields. *Journal of The Royal Society Interface*, 12(103), 20141103.

WILDLIFE , WIRELESS AND NON IONIZING ELECTROMAGNETIC FIELDS

SCIENTIFIC RESEARCH

POLLINATORS

A. Lázaro, A. Chroni, T. Tscheulin, J. Devalez, C. Matsoukas, & T. Petanidou. (2016). Electromagnetic radiation of mobile telecommunication antennas affects the abundance and composition of wild pollinators. *Journal of Insect Conservation*, 20(2), 315–324. <https://doi.org/10.1007/s10841-016-9868-8>

Adelaja, O. J., Ande, A. T., Abdulraheem, G. D., Oluwakorode, I. A., Oladipo, O. A., & Oluwajobi, A. O. (2021). Distribution, diversity and abundance of some insects around a telecommunication mast in Ilorin, Kwara State, Nigeria. *Bulletin of the National Research Centre*, 45(1), 222.

Balmori A. (2021) Electromagnetic radiation as an emerging driver factor for the decline of insects. *Science of the Total Environment*. 767: 144913

Borre, E. D., Joseph, W., Aminzadeh, R., Müller, P., Boone, M. N., Josipovic, I., Hashemizadeh, S., Kuster, N., Kühn, S., & Thielens, A. (2021). Radio-frequency exposure of the yellow fever mosquito (*A. aegypti*) from 2 to 240 GHz. *PLOS Computational Biology*, 17(10), e1009460.

Favre, D. (2011). Mobile phone-induced honeybee worker piping. *Apidologie*, 42(3), 270–279.

Fedele, G., Edwards, M. D., Bhutani, S., Hares, J. M., Murbach, M., Green, E. W., Dissel, S., Hastings, M. H., Rosato, E., & Kyriacou, C. P. (2014). Genetic analysis of circadian responses to low frequency electromagnetic fields in *Drosophila melanogaster*. *PLoS Genetics*, 10(12), e1004804.

Lee, K.-S., Choi, J.-S., Hong, S.-Y., Son, T.-H., & Yu, K. (2008). Mobile phone electromagnetic radiation activates MAPK signaling and regulates viability in *Drosophila*. *Bioelectromagnetics*, 29(5), 371–379.

Li, S.-S., Zhang, Z.-Y., Yang, C.-J., Lian, H.-Y., & Cai, P. (2013). Gene expression and reproductive abilities of male *Drosophila melanogaster* subjected to ELF-EMF exposure. *Mutation Research. Genetic Toxicology and Environmental Mutagenesis*, 758(1–2), 95–103.

Lopatina, N. G., Zachepilo, T. G., Kamyshev, N. G., Dyuzhikova, N. A., & Serov, I. N. (2019). Effect of Non-Ionizing Electromagnetic Radiation on Behavior of the Honeybee, *Apis mellifera* L. (Hymenoptera, Apidae). *Entomological Review*, 99(1), 24–29.

Lupi, D., Palamara Mesiano, M., Adani, A., Benocci, R., Giacchini, R., Parenti, P., Zambon, G., Lavazza, A., Boniotti, M. B., Bassi, S., Colombo, M., & Tremolada, P. (2021a). Combined Effects of Pesticides and Electromagnetic-Fields on Honeybees: Multi-Stress Exposure. *Insects*, 12(8), 716.

erator affects the sleep of *Drosophila melanogaster*. *Environmental Pollution (Barking, Essex: 1987)*, 283, 117087.

WILDLIFE , WIRELESS AND NON IONIZING ELECTROMAGNETIC FIELDS

SCIENTIFIC RESEARCH

Manta, A. K., Papadopoulou, D., Polyzos, A. P., Fragopoulou, A. F., Skouroliakou, A. S., Thanos, D., Stravopodis, D. J., & Margaritis, L. H. (2017). Mobile-phone radiation-induced perturbation of gene-expression profiling, redox equilibrium and sporadic-apoptosis control in the ovary of *Drosophila melanogaster*. *Fly*, 11(2), 75-95.

Mahmoud EA and Gabarty A (2021) "Impact of Electromagnetic Radiation on Honey Stomach Ultrastructure and the Body Chemical Element Composition of *Apis mellifera*." *African Entomology* 29(1), 32-41, (23 March).

Molina-Montenegro MA, Acuña-Rodríguez IS, Ballesteros GI, Baldelomar M, Torres-Díaz C, Broitman BR, Vázquez DP. (2023) Electromagnetic fields disrupt the pollination service by honeybees. *Sci Adv*. May 12;9(19)

Migdał, P., Berbeć, E., Bieńkowski, P., Plotnik, M., Murawska, A., & Latarowski, K. (2022b). Exposure to Magnetic Fields Changes the Behavioral Pattern in Honeybees (*Apis mellifera* L.) under Laboratory Conditions. *Animals: An Open Access Journal from MDPI*, 12(7), 855.

Odemer, R., & Odemer, F. (2019). Effects of radiofrequency electromagnetic radiation (RF-EMF) on honey bee queen development and mating success. *Science of The Total Environment*, 661, 553-562.

Santhosh Kumar, S. (2018). Colony Collapse Disorder (CCD) in Honey Bees Caused by EMF Radiation. *Bioinformation*, 14(9), 421-424.

Thielens, A., Bell, D., Mortimore, D. B., Greco, M. K., Martens, L., & Joseph, W. (2018). Exposure of Insects to Radio-Frequency Electromagnetic Fields from 2 to 120 GHz. *Scientific Reports*, 8(1), 3924.

Thielens A, Greco MK, Verloock L, Martens L, Joseph W. Radio-Frequency Electromagnetic Field Exposure of Western Honey Bees. *Scientific Reports*. 2020 Jan 16;10(1):461.

Wang, Y., Jiang, Z., Zhang, L., Zhang, Z., Liao, Y., & Cai, P. (2022b). 3.5-GHz radiofrequency electromagnetic radiation promotes the development of *Drosophila melanogaster*. *Environmental Pollution (Barking, Essex: 1987)*, 294, 118646.

Wang, Y., Zhang, H., Zhang, Z., Sun, B., Tang, C., Zhang, L., Jiang, Z., Ding, B., Liao, Y., & Cai, P. (2021). Simulated mobile communication frequencies (3.5 GHz) emitted by a signal generator affects the sleep of *Drosophila melanogaster*. *Environmental Pollution (Barking, Essex: 1987)*, 283, 117087.

AMPHIBIAN

Balmori A. (2010). Mobile phone mast effects on common frog (*Rana temporaria*) tadpoles: the city turned into a laboratory. *Electromagn Biol Med*. Jun;29 (1-2): 31-5.

Balmori, A. (2006). The incidence of electromagnetic pollution on the amphibian decline: Is this an important piece of the puzzle? *Toxicological & Environmental Chemistry*, 88(2), 287-299.

WILDLIFE , WIRELESS AND NON IONIZING ELECTROMAGNETIC FIELDS

SCIENTIFIC RESEARCH

PLANTS

Halgamuge, M. N. (2017). [Review: Weak radiofrequency radiation exposure from mobile phone radiation on plants.](#) *Electromagnetic Biology and Medicine*, 36(2), 213–235.

Halgamuge, M. N., Yak, S. K., & Eberhardt, J. L. (2015). [Reduced growth of soybean seedlings after exposure to weak microwave radiation from GSM 900 mobile phone and base station.](#) *Bioelectromagnetics*, 36(2), 87–95.

Haggerty, K. (2010). [Adverse Influence of Radio Frequency Background on Trembling Aspen Seedlings: Preliminary Observations.](#) *International Journal of Forestry Research*, 836278.

Kaur, S., Vian, A., Chandel, S., Singh, D. H., Batish, D., & Kohli, R. (2021). [Sensitivity of plants to high frequency electromagnetic radiation: Cellular mechanisms and morphological changes.](#) *Reviews in Environmental Science and Bio/Technology*, 20.

Soran, M.-L., Stan, M., Niinemets, Ü., & Copolovici, L. (2014). [Influence of microwave frequency electromagnetic radiation on terpene emission and content in aromatic plants.](#) *Journal of Plant Physiology*, 171(15), 1436–1443.

Stefi, A. L., Margaritis, L. H., & Christodoulakis, N. S. (2016). [The effect of the non ionizing radiation on cultivated plants of Arabidopsis thaliana \(Col.\).](#) *Flora*, 223, 114–120.

Waldmann-Selsam, C., Balmori-de la Puente, A., Breunig, H., & Balmori, A. (2016). [Radiofrequency radiation injures trees around mobile phone base stations.](#) *Science of The Total Environment*, 572, 554–569.

Zhong, Z., Wang, X., Yin, X., Tian, J., & Komatsu, S. (2021). [Morphophysiological and Proteomic Responses on Plants of Irradiation with Electromagnetic Waves.](#) *International Journal of Molecular Sciences*, 22(22), Article 22.

MARINE

Hutchison, Z. L., Gill, A. B., Sigray, P., He, H., & King, J. W. (2020). [Anthropogenic electromagnetic fields \(EMF\) influence the behaviour of bottom-dwelling marine species.](#) *Scientific Reports*, 10(1), 4219.

Scott, K., Harsanyi, P., Easton, B. A. A., Piper, A. J. R., Rochas, C. M. V., & Lyndon, A. R. (2021). [Exposure to Electromagnetic Fields \(EMF\) from Submarine Power Cables Can Trigger Strength-Dependent Behavioural and Physiological Responses in Edible Crab, Cancer pagurus \(L.\).](#) *Journal of Marine Science and Engineering*, 9(7), Article 7.

WILDLIFE , WIRELESS AND NON IONIZING ELECTROMAGNETIC FIELDS

SCIENTIFIC RESEARCH

PLANTS

Halgamuge, M. N. (2017). [Review: Weak radiofrequency radiation exposure from mobile phone radiation on plants.](#) *Electromagnetic Biology and Medicine*, 36(2), 213–235.

Halgamuge, M. N., Yak, S. K., & Eberhardt, J. L. (2015). [Reduced growth of soybean seedlings after exposure to weak microwave radiation from GSM 900 mobile phone and base station.](#) *Bioelectromagnetics*, 36(2), 87–95.

Haggerty, K. (2010). [Adverse Influence of Radio Frequency Background on Trembling Aspen Seedlings: Preliminary Observations.](#) *International Journal of Forestry Research*, 836278.

Kaur, S., Vian, A., Chandel, S., Singh, D. H., Batish, D., & Kohli, R. (2021). [Sensitivity of plants to high frequency electromagnetic radiation: Cellular mechanisms and morphological changes.](#) *Reviews in Environmental Science and Bio/Technology*, 20.

Soran, M.-L., Stan, M., Niinemets, Ü., & Copolovici, L. (2014). [Influence of microwave frequency electromagnetic radiation on terpene emission and content in aromatic plants.](#) *Journal of Plant Physiology*, 171(15), 1436–1443.

Stefi, A. L., Margaritis, L. H., & Christodoulakis, N. S. (2016). [The effect of the non ionizing radiation on cultivated plants of Arabidopsis thaliana \(Col.\).](#) *Flora*, 223, 114–120.

Waldmann-Selsam, C., Balmori-de la Puente, A., Breunig, H., & Balmori, A. (2016). [Radiofrequency radiation injures trees around mobile phone base stations.](#) *Science of The Total Environment*, 572, 554–569.

Zhong, Z., Wang, X., Yin, X., Tian, J., & Komatsu, S. (2021). [Morphophysiological and Proteomic Responses on Plants of Irradiation with Electromagnetic Waves.](#) *International Journal of Molecular Sciences*, 22(22), Article 22.

Additional Studies To Know

Balmori A. (2010). [Mobile phone mast effects on common frog \(Rana temporaria\) tadpoles: the city turned into a laboratory.](#) *Electromagn Biol Med.* Jun;29 (1-2): 31-5.

Hutchison, Z. L., Gill, A. B., Sigray, P., He, H., & King, J. W. (2020). [Anthropogenic electromagnetic fields \(EMF\) influence the behaviour of bottom-dwelling marine species.](#) *Scientific Reports*, 10(1), 4219.

Scott, K., Harsanyi, P., Easton, B. A. A., Piper, A. J. R., Rochas, C. M. V., & Lyndon, A. R. (2021). [Exposure to Electromagnetic Fields \(EMF\) from Submarine Power Cables Can Trigger Strength-Dependent Behavioural and Physiological Responses in Edible Crab, Cancer pagurus \(L.\).](#) *Journal of Marine Science and Engineering*, 9(7), Article



American Academy of Pediatrics Webpage Excerpts

Electromagnetic Fields: A Hazard to Your Health?

In recent years, concern has increased about exposure to radio frequency electromagnetic radiation emitted from cell phones and phone station antennae. An Egyptian study confirmed concerns that living nearby mobile phone base stations increased the risk for developing:

- Headaches
- Memory problems
- Dizziness
- Depression
- Sleep problems

Short-term exposure to these fields in experimental studies have not always shown negative effects, but this does not rule out cumulative damage from these fields, so larger studies over longer periods are needed to help understand who is at risk. In large studies, an association has been observed between symptoms and exposure to these fields in the everyday environment.

Last Updated 12/28/2012

Source American Academy of Pediatrics (Copyright © 2012)



Environmental Pollution

journal homepage: www.elsevier.com/locate/envpol



Thermal and non-thermal health effects of low intensity non-ionizing radiation: An international perspective[☆]

Dominique Belpomme^{a, b, 1}, Lennart Hardell^{a, c, 1, 2}, Igor Belyaev^{a, d, e, 1}, Ernesto Burgio^{a, f}, David O. Carpenter^{a, g, h, *, 1}

^a European Cancer Environment Research Institute, Brussels, Belgium

^b Paris V University Hospital, Paris, France

^c Department of Oncology, Orebro University Hospital, Faculty of Medicine, Orebro, Sweden

^d Department of Radiobiology, Cancer Research Institute, Biomedical Research Center, Slovak Academy of Science, Bratislava, Slovak Republic

^e Laboratory of Radiobiology, Institute of General Physics, Russian Academy of Science, Moscow, Russian Federation

^f Istituto Scientifico Biomedico Euro Mediterraneo, Mesagne, Italy

^g Institute for Health and the Environment, University at Albany, Albany, NY, USA

^h Child Health Research Centre, The University of Queensland, Faculty of Medicine, Brisbane, Australia



ARTICLE INFO

Article history:

Received 6 April 2018

Received in revised form

31 May 2018

Accepted 4 July 2018

Available online 6 July 2018

ABSTRACT

Exposure to low frequency and radiofrequency electromagnetic fields at low intensities poses a significant health hazard that has not been adequately addressed by national and international organizations such as the World Health Organization. There is strong evidence that excessive exposure to mobile phone-frequencies over long periods of time increases the risk of brain cancer both in humans and animals. The mechanism(s) responsible include induction of reactive oxygen species, gene expression alteration and DNA damage through both epigenetic and genetic processes. *In vivo* and *in vitro* studies demonstrate adverse effects on male and female reproduction, almost certainly due to generation of reactive oxygen species. There is increasing evidence the exposures can result in neurobehavioral decrements and that some individuals develop a syndrome of "electro-hypersensitivity" or "microwave illness", which is one of several syndromes commonly categorized as "idiopathic environmental intolerance". While the symptoms are non-specific, new biochemical indicators and imaging techniques allow diagnosis that excludes the symptoms as being only psychosomatic. Unfortunately standards set by most national and international bodies are not protective of human health. This is a particular concern in children, given the rapid expansion of use of wireless technologies, the greater susceptibility of the developing nervous system, the hyperconductivity of their brain tissue, the greater penetration of radiofrequency radiation relative to head size and their potential for a longer lifetime exposure.

Biological effects from exposure to electromagnetic radiation emitted by cell tower base stations and other antenna arrays

B. Blake Levitt and Henry Lai

Abstract: The siting of cellular phone base stations and other cellular infrastructure such as roof-mounted antenna arrays, especially in residential neighborhoods, is a contentious subject in land-use regulation. Local resistance from nearby residents and landowners is often based on fears of adverse health effects despite reassurances from telecommunications service providers that international exposure standards will be followed. Both anecdotal reports and some epidemiology studies have found headaches, skin rashes, sleep disturbances, depression, decreased libido, increased rates of suicide, concentration problems, dizziness, memory changes, increased risk of cancer, tremors, and other neurophysiological effects in populations near base stations. The objective of this paper is to review the existing studies of people living or working near cellular infrastructure and other pertinent studies that could apply to long-term, low-level radiofrequency radiation (RFR) exposures. While specific epidemiological research in this area is sparse and contradictory, and such exposures are difficult to quantify given the increasing background levels of RFR from myriad personal consumer products, some research does exist to warrant caution in infrastructure siting. Further epidemiology research that takes total ambient RFR exposures into consideration is warranted. Symptoms reported today may be classic microwave sickness, first described in 1978. Non-ionizing electromagnetic fields are among the fastest growing forms of environmental pollution. Some extrapolations can be made from research other than epidemiology regarding biological effects from exposures at levels far below current exposure guidelines.



Clinical Biochemistry

Volume 45, Issues 1–2, January 2012, Pages 157–161



Case Report

How does long term exposure to base stations and mobile phones affect human hormone profiles?

Emad F. Eskander , Selim F. Estefan, Ahmed A. Abd-Rabou

Objectives

This study is concerned with assessing the role of exposure to radio frequency radiation (RFR) emitted either from mobiles or base stations and its relations v human's hormone profiles.

Results

This study showed significant decrease in volunteers' ACTH, cortisol, thyroid hormones, prolactin for young females, and testosterone levels.



Very high radiofrequency radiation at Skeppsbron in Stockholm, Sweden from mobile phone base station antennas positioned close to pedestrians' heads

Tarmo Koppel^a, Mikko Ahonen^b, Michael Carlberg^c, Lennart Hardell^{c,*}

^a Tallinn University of Technology, SOCS53 Ehitajate tee 5, 19086, Tallinn, Estonia

^b Päiväliite, Päivölkantie 52, Tarvita, 37770, Finland

^c The Environment and Cancer Research Foundation, Studievägen 35, SE 702 17, Örebro, Sweden



Fig. 3. Street view on the Skeppsbron street with some of the mobile phone base station antennas pointed out with a circle; note the low placement of the antennas, where microwaves irradiate the pedestrian at close range.

ABSTRACT

In urban environment there is a constant increase of public exposure to radiofrequency electromagnetic fields from mobile phone base stations. With the placement of mobile phone base station antennas radiofrequency hotspots emerge. This study investigates an area at Skeppsbron street in Stockholm, Sweden with an aggregation of base station antennas placed at low level close to pedestrians' heads. Detailed spatial distribution measurements were performed with 1) a radiofrequency broadband analyzer and 2) a portable exposimeter. The results display a greatly uneven distribution of the radiofrequency field with hotspots. The highest spatial average across all quadrat cells was 12.1 V m^{-1} (388 mW m^{-2}), whereas the maximum recorded reading from the entire area was 31.6 V m^{-1} (2648 mW m^{-2}). Exposimeter measurements show that the majority of exposure is due to mobile phone downlink bands. Most dominant are 2600 and 2100 MHz bands used by 4G and 3G mobile phone services, respectively. The average radiofrequency radiation values from the earlier studies show that the level of ambient RF radiation exposure in Stockholm is increasing. This study concluded that mobile phone base station antennas at Skeppsbron, Stockholm are examples of poor radiofrequency infrastructure design which brings upon highly elevated exposure levels to popular seaside promenade and a busy traffic street.

Studies from recent decades have shown elevated health risk under long term exposure to such highly elevated radiofrequency fields.

A review by [Khurana et al. \(2010\)](#) found in 80% of the available studies neurobehavioral symptoms or cancer in populations living at distances <500 m from base stations ([Khurana et al., 2010](#)). In another review exposure from base stations and other antenna arrays showed changes in immunological and reproductive systems as well as DNA double strand breaks, influence on calcium movement in the heart and increased proliferation rates in human astrocytoma cancer cells ([Levitt and Lai, 2010](#)).

When a GSM 900 MHz base station was installed in the village Rimbach in Germany it had an influence on the neurotransmitters adrenaline, noradrenaline, dopamine and phenylethylamine ([Buchner and Eger, 2011](#)). Influence on cortisol and thyroid hormones in people living near base stations was shown in other studies ([Augner et al., 2010](#); [Eskander et al., 2012](#)).

[Dode et al. \(2011\)](#) compared base station (BS) clusters and cases of deaths by neoplasia in the Belo Horizonte municipality, Minas Gerais state, Brazil, from 1996 to 2006. In their study largest electric field was 12.4 V m^{-1} and the smallest was 0.4 V m^{-1} . They found cancer-related death rates be higher close to base stations. This finding confirmed earlier findings by [Eger \(2004\)](#).

In a study from India, genetic damage using the single cell gel electrophoresis (comet) assay was assessed in peripheral blood leukocytes of individuals residing in the vicinity of a mobile phone base station and comparing it to that in healthy controls. Genetic damage parameters of DNA migration length, damage frequency, and damage index were significantly ($p < 0.001$) elevated in the sample group compared to respective values in healthy controls ([Gandhi et al., 2014](#)).

The effect of RF radiation among 20 subjects living close to mobile phone base station compared with 20 subjects living with a distance of about 1 km was studied ([Singh et al., 2016](#)). The authors concluded that: "It was unveiled that a majority of the subjects who were residing near the mobile base station complained of sleep disturbances, headache, dizziness, irritability, concentration difficulties, and hypertension. A majority of the study subjects had significantly lesser stimulated salivary secretion ($p < 0.01$) as compared to the control subjects."

[Zothansiana et al. \(2017\)](#) in India inspected DNA damage and antioxidant status in cultured human peripheral blood lymphocytes (HPBLs) of individuals residing in the vicinity of mobile phone base stations and compared it with healthy controls living further away. Analyses of data from the exposed group ($n = 40$), residing within a perimeter of 80 m of mobile base stations, showed statistically significantly ($p < 0.0001$) higher frequency of micronuclei when compared to the control group, residing 300 m away from the mobile base station.

The Ramazzini Institute findings ([Falcioni et al., 2018](#)) are supported by the results in the USNTP study on rats and mice exposed to RF radiation ([National Toxicology Program, 2018a, 2018b](#)). A clear evidence of increased incidence of heart Schwannoma and some evidence of glioma and tumours in the adreanal medulla in male rats was found according to the expert panel, for further discussion see [Hardell and Carlberg \(2019\)](#).

The study concluded that Skeppsbron street mobile phone base station antennas are examples of a poor radiofrequency infrastructure design with mobile phone base station antennas positioned into close range to the general public which brings upon high exposure levels. Given the low placement of the antennas (height from the street floor), the highest exposure was often registered at pedestrian head level. Given that head is one of most vulnerable parts of the body, these placements by mobile telephony service providers put pedestrians into unnecessary risk. Position of these antennas, can pose a health risk to people at close range. This is especially critical for people at particular risk, including persons with medical implants, pregnant women or chronically ill persons.

Based on the latest scientific literature regarding RF exposure and adverse health effects, this study recommends repositioning such base station antennas to areas away from the nearby inhabitants, workers and the general public. Alternatively, very low power antennas may also be considered to reduce the exposure. Occupational exposure of people



Low Intensity Electromagnetic Fields Act *via* Voltage-Gated Calcium Channel (VGCC) Activation to Cause Very Early Onset Alzheimer's Disease: 18 Distinct Types of Evidence

Martin L. Pall^{1,*}

¹Professor Emeritus of Biochemistry & Basic Medical Sciences, |Washington State University, Current Address: 638 NE 41st Ave., Portland, OR 97232, USA

ARTICLE HISTORY

Received: October 04, 2021
Revised: December 22, 2021
Accepted: December 31, 2021

DOI:
10.2174/1567205019666220202114510



CrossMark


This is an Open Access article published under CC BY 4.0
<https://creativecommons.org/licenses/by/4.0/legalcode>

Abstract: Electronically generated electromagnetic fields (EMFs), including those used in wireless communication such as cell phones, Wi-Fi and smart meters, are coherent, producing very high electric and magnetic forces, which act on the voltage sensor of voltage-gated calcium channels to produce increases in intracellular calcium $[Ca^{2+}]_i$. The calcium hypothesis of Alzheimer's disease (AD) has shown that each of the important AD-specific and nonspecific causal elements is produced by excessive $[Ca^{2+}]_i$. $[Ca^{2+}]_i$ acts in AD *via* excessive calcium signaling and the peroxynitrite/oxidative stress/inflammation pathway, which are each elevated by EMFs. An apparent vicious cycle in AD involves amyloid-beta protein ($A\beta$) and $[Ca^{2+}]_i$. Three types of epidemiology suggest EMF causation of AD, including early onset AD. Extensive animal model studies show that low intensity EMFs cause neurodegeneration, including AD, with AD animals having elevated levels of $A\beta$, amyloid precursor protein and BACE1. Rats exposed to pulsed EMFs every day are reported to develop universal or near universal very early onset neurodegeneration, including AD; these findings are superficially similar to humans with digital dementia. EMFs producing modest increases in $[Ca^{2+}]_i$ can also produce protective, therapeutic effects. The therapeutic pathway and peroxynitrite pathway inhibit each other. A summary of 18 different findings is provided, which collectively provide powerful evidence for EMF causation of AD. The author is concerned that smarter, more highly pulsed "smart" wireless communication may cause widespread very, very early onset AD in human populations.

ELECTROMAGNETIC BIOLOGY AND MEDICINE
2022, VOL. 41, NO. 2, 230-255
<https://doi.org/10.1080/15368378.2022.2065683>



Taylor & Francis
Taylor & Francis Group



The roles of intensity, exposure duration, and modulation on the biological effects of radiofrequency radiation and exposure guidelines

Henry Lai^a and B. Blake Levitt^b

^aDepartment of Bioengineering, University of Washington, Seattle, WA, USA; ^bNew Preston, CT, USA

ABSTRACT

In this paper, we review the literature on three important exposure metrics that are inadequately represented in most major radiofrequency radiation (RFR) exposure guidelines today: intensity, exposure duration, and signal modulation. Exposure intensity produces unpredictable effects as demonstrated by nonlinear effects. This is most likely caused by the biological system's ability to adjust and compensate but could lead to eventual biomic breakdown after prolonged exposure. A review of 112 low-intensity studies reveals that biological effects of RFR could occur at a median specific absorption rate of 0.0165 W/kg. Intensity and exposure duration interact since the dose of energy absorbed is the product of intensity and time. The result is that RFR behaves like a biological "stressor" capable of affecting numerous living systems. In addition to intensity and duration, man-made RFR is generally modulated to allow information to be encrypted. The effects of modulation on biological functions are not well understood. Four types of modulation outcomes are discussed. In addition, it is invalid to make direct comparisons between thermal energy and radiofrequency electromagnetic energy. Research data indicate that electromagnetic energy is more biologically potent in causing effects than thermal changes. The two likely function through different mechanisms. As such, any current RFR exposure guidelines based on acute continuous-wave exposure are inadequate for health protection.

ARTICLE HISTORY

Received 15 February 2022
Accepted 1 April 2022

KEYWORDS

Radiofrequency radiation (RFR); intensity; duration of exposure; modulation; specific absorption rate (SAR); biological effects

BENTHAM
SCIENCE

Low Intensity Electromagnetic Fields Act *via* Voltage-Gated Calcium Channel (VGCC) Activation to Cause Very Early Onset Alzheimer's Disease: 18 Distinct Types of Evidence

Martin L. Pall^{1,*}

¹Professor Emeritus of Biochemistry & Basic Medical Sciences, |Washington State University, Current Address: 638 NE 41st Ave., Portland, OR 97232, USA

ARTICLE HISTORY

Received: October 04, 2021
Revised: December 22, 2021
Accepted: December 31, 2021

DOI:
10.2174/15672050196662202114510



CrossMark


This is an Open Access article published under CC BY 4.0
<https://creativecommons.org/licenses/by/4.0/legalcode>

Abstract: Electronically generated electromagnetic fields (EMFs), including those used in wireless communication such as cell phones, Wi-Fi and smart meters, are coherent, producing very high electric and magnetic forces, which act on the voltage sensor of voltage-gated calcium channels to produce increases in intracellular calcium $[Ca^{2+}]_i$. The calcium hypothesis of Alzheimer's disease (AD) has shown that each of the important AD-specific and nonspecific causal elements is produced by excessive $[Ca^{2+}]_i$. $[Ca^{2+}]_i$ acts in AD *via* excessive calcium signaling and the peroxynitrite/oxidative stress/inflammation pathway, which are each elevated by EMFs. An apparent vicious cycle in AD involves amyloid-beta protein ($A\beta$) and $[Ca^{2+}]_i$. Three types of epidemiology suggest EMF causation of AD, including early onset AD. Extensive animal model studies show that low intensity EMFs cause neurodegeneration, including AD, with AD animals having elevated levels of $A\beta$, amyloid precursor protein and BACE1. Rats exposed to pulsed EMFs every day are reported to develop universal or near universal very early onset neurodegeneration, including AD; these findings are superficially similar to humans with digital dementia. EMFs producing modest increases in $[Ca^{2+}]_i$ can also produce protective, therapeutic effects. The therapeutic pathway and peroxynitrite pathway inhibit each other. A summary of 18 different findings is provided, which collectively provide powerful evidence for EMF causation of AD. The author is concerned that smarter, more highly pulsed "smart" wireless communication may cause widespread very, very early onset AD in human populations.

ELECTROMAGNETIC BIOLOGY AND MEDICINE
<https://doi.org/10.1080/15368378.2021.1881866>

Taylor & Francis
Taylor & Francis Group

REVIEW



Genetic effects of non-ionizing electromagnetic fields

Henry Lai

Department of Bioengineering, University of Washington, Seattle, WA, USA

ARTICLE HISTORY

Received 22 September 2020
Accepted 13 December 2020

KEYWORDS



Radiofrequency radiation; static/extremely low frequency EMF; genetic effects; genotoxicity; gene expression

ABSTRACT

This is a review of the research on the genetic effects of non-ionizing electromagnetic field (EMF), mainly on radiofrequency radiation (RFR) and static and extremely low frequency EMF (ELF-EMF). The majority of the studies are on genotoxicity (e.g., DNA damage, chromatin conformation changes, etc.) and gene expression. Genetic effects of EMF depend on various factors, including field parameters and characteristics (frequency, intensity, wave-shape), cell type, and exposure duration. The types of gene expression affected (e.g., genes involved in cell cycle arrest, apoptosis and stress responses, heat-shock proteins) are consistent with the findings that EMF causes genetic damages. Many studies reported effects in cells and animals after exposure to EMF at intensities similar to those in the public and occupational environments. The mechanisms by which effects are induced by EMF are basically unknown. Involvement of free radicals is a likely possibility. EMF also interacts synergistically with different entities on genetic functions. Interactions, particularly with chemotherapeutic compounds, raise the possibility of using EMF as an adjuvant for cancer treatment to increase the efficacy and decrease side effects of traditional chemotherapeutic drugs. Other data, such as adaptive effects and mitotic spindle aberrations after EMF exposure, further support the notion that EMF causes genetic effects in living organisms.

Article

The Effect of Continuous Low-Intensity Exposure to Electromagnetic Fields from Radio Base Stations to Cancer Mortality in Brazil

Nádia Cristina Pinheiro Rodrigues ^{1,2,*} , Adilza Condessa Dode ³, Mônica Kramer de Noronha Andrade ¹ , Gisele O'Dwyer ¹, Denise Leite Maia Monteiro ⁴, Inês Nascimento Carvalho Reis ¹, Roberto Pinheiro Rodrigues ^{5,6}, Vera Cecília Frossard ¹ and Valéria Teresa Saraiva Lino ¹

Abstract: Background: this study aims to estimate the rate of death by cancer as a result of Radio Base Station (RBS) radiofrequency exposure, especially for breast, cervix, lung, and esophagus cancers. Methods: we collected information on the number of deaths by cancer, gender, age group, gross domestic product per capita, death year, and the amount of exposure over a lifetime. We investigated all cancer types and some specific types (breast, cervix, lung, and esophagus cancers). Results: in capitals where RBS radiofrequency exposure was higher than 2000/antennas-year, the average mortality rate was 112/100,000 for all cancers. The adjusted analysis showed that, the higher the exposure to RBS radiofrequency, the higher cancer mortality was. The highest adjusted risk was observed for cervix cancer (rate ratio = 2.18). The spatial analysis showed that the highest RBS radiofrequency exposure was observed in a city in southern Brazil that also showed the highest mortality rate for all types of cancer and specifically for lung and breast cancer. Conclusion: the balance of our results indicates that exposure to radiofrequency electromagnetic fields from RBS increases the rate of death for all types of cancer.



Contents lists available at ScienceDirect

Environmental Research

journal homepage: www.elsevier.com/locate/envres



Report of final results regarding brain and heart tumors in Sprague-Dawley rats exposed from prenatal life until natural death to mobile phone radiofrequency field representative of a 1.8 GHz GSM base station environmental emission

ABSTRACT

Background: In 2011, IARC classified radiofrequency radiation (RFR) as possible human carcinogen (Group 2B). According to IARC, animals studies, as well as epidemiological ones, showed limited evidence of carcinogenicity. In 2016, the NTP published the first results of its long-term bioassays on near field RFR, reporting increased incidence of malignant glial tumors of the brain and heart Schwannoma in rats exposed to GSM – and CDMA – modulated cell phone RFR. The tumors observed in the NTP study are of the type similar to the ones observed in some epidemiological studies of cell phone users.

Objectives: The Ramazzini Institute (RI) performed a life-span carcinogenic study on Sprague-Dawley rats to evaluate the carcinogenic effects of RFR in the situation of far field, reproducing the environmental exposure to RFR generated by 1.8 GHz GSM antenna of the radio base stations of mobile phone. This is the largest long-term study ever performed in rats on the health effects of RFR, including 2448 animals. In this article, we reported the final results regarding brain and heart tumors.

Methods: Male and female Sprague-Dawley rats were exposed from prenatal life until natural death to a 1.8 GHz GSM far field of 0, 5, 25, 50 V/m with a whole-body exposure for 19 h/day.

Results: A statistically significant increase in the incidence of heart Schwannomas was observed in treated male rats at the highest dose (50 V/m). Furthermore, an increase in the incidence of heart Schwann cells hyperplasia was observed in treated male and female rats at the highest dose (50 V/m), although this was not statistically significant. An increase in the incidence of malignant glial tumors was observed in treated female rats at the highest dose (50 V/m), although not statistically significant.

Conclusions: The RI findings on far field exposure to RFR are consistent with and reinforce the results of the NTP study on near field exposure, as both reported an increase in the incidence of tumors of the brain and heart in RFR-exposed Sprague-Dawley rats. These tumors are of the same histotype of those observed in some epidemiological studies on cell phone users. These experimental studies provide sufficient evidence to call for the re-evaluation of IARC conclusions regarding the carcinogenic potential of RFR in humans.