



Chris Ketchen
Town Manager
6 Walker Street
Lenox, MA 01240

Dear Lenox Town Manager Chris Ketchen,

We write to you regarding the placement of cell towers, “small cell” and wireless facilities. Cell antennas and cell towers should not be placed near schools and homes. Safety is not assured even if the cell tower is compliant with FCC limits.

On August 13, 2021, the United States Court of Appeals for the District of Columbia Circuit [ruled in our case](#) against the FCC that the decision by the Federal Communications Commission (FCC) not to update its 1996 safety limits for human exposure to wireless radiation (which includes cell tower emissions) was “arbitrary and capricious.” One of the most important aspects of the court decision was that the court found the FCC did not adequately explain why it ignored scientific evidence on impacts from long term wireless radiation exposure, especially in regards to children, who the American Academy of Pediatrics states are more vulnerable to wireless radiation. The court ordered the FCC to examine the record evidence regarding long term exposure to children. So far, the FCC has not responded. Thus, this landmark [ruling](#) highlights how no federal health agency has reviewed the full body of current research to ensure the current human exposure limits of the FCC are adequately protective.

Extensive published scientific evidence indicates that radiofrequency radiation *at levels far below FCC limits* can cause [cancer](#), [increased oxidative stress](#), [genetic damage](#), structural and functional changes of the [reproductive system](#), [memory deficits](#), [behavioral problems](#), and [neurological impacts](#). We consider radiofrequency radiation (RFR) to be a human carcinogen based on the [current body](#) of evidence.

Please see our latest publication entitled “[Wireless technologies, non-ionizing electromagnetic fields and children: Identifying and reducing health risks](#)” published in *Current Problems in Pediatric and Adolescent Health Care* regarding the need to mitigate risk to children.

The exposure limits of the US Federal Communications Commission are totally outdated and do not protect the health of the public, especially not the health of children. The Los Angeles School District has banned cell towers on their District’s school grounds and also [set an RF limit at 10,000 less than FCC](#)

[limits](#) stating that, “It is believed that a more conservative level is necessary to protect children, who represent a potentially vulnerable and sensitive population.”

Children are more vulnerable to RF radiation, just as they are to other environmental exposures. Even very low exposures to children can have [serious impacts](#) later in life because their nervous and immune systems are rapidly developing. Children absorb higher levels of RF radiation deeper into their brains and bodies because they have thinner skulls, a higher water content in their tissues. Children will have a longer lifetime of higher exposures, compared to adults today.

The [American Academy of Pediatrics states](#):

“In recent years, concern has increased about exposure to radio frequency (RF) electromagnetic radiation emitted from cell phones and phone station antennas. An Egyptian study confirmed concerns that living nearby mobile phone base stations increased the risk for developing:

- Headaches
- Memory problems
- Dizziness
- Depression
- Sleep problems

Short-term exposure to these fields in experimental studies have not always shown negative effects, but this does not rule out cumulative damage from these fields, so larger studies over longer periods are needed to help understand who is at risk. In large studies, an association has been observed between symptoms and exposure to these fields in the everyday environment.”

At this time we have not identified a safe level of exposure. Although radiation levels decrease as you increase your distance from a particular antenna/tower, the reality is that adding a tower or base station to a community will definitely *increase* the radiation exposure in that area near the antennas and within the surrounding coverage area.

We recommend policies to reduce human exposure to RFR, especially for children. Schools are where children spend the majority of their daytime hours. Therefore we strongly recommend against installing cell towers near schools, daycares, parks, homes, or hospitals.

Recent research on people living near cell antennas has found increases in molecular markers in the blood that predict cancer. This study evaluated effects in the human blood of individuals living near mobile phone base stations (for study purposes, they chose a distance of 80 meters) compared with healthy controls living more than 300 meters from a base station. The study measured higher RFR levels in the homes of people living in homes within 80 meters from the cell antennas (documenting the impact of increased RFR radiation from the antenna installations) and found statistically significant differences in their blood. The group living closer to the antennas had statistically significant higher frequency of micronuclei and a rise in lipid peroxidation in their blood; these changes are considered biomarkers predictive of cancer ([Zothansiama et al. 2017](#)).

A review paper entitled “[Limiting liability with positioning to minimize negative health effects of cellular phone towers](#)” reviewed the “large and growing body of evidence that human exposure to RFR from cellular phone base stations causes negative health effects.” The authors recommend restricting antennas near home and within 500 meters of schools and hospitals to protect companies from future liability ([Pearce 2020](#)).

Please note the following scientific publications regarding cell towers and cell phone radiation:

- European Parliament requested a research report “[Health Impact of 5G](#)” which was released in July 2021 and concluded that commonly used RFR frequencies (450 to 6000 MHz) are probably carcinogenic for humans and clearly affect male fertility with possible adverse effects on the development of embryos, fetuses and newborns.
- A review entitled “[Evidence for a health risk by RF on humans living around mobile phone base stations: From radiofrequency sickness to cancer](#)” reviewed the existing scientific literature and found radiofrequency sickness, cancer and changes in biochemical parameters ([Balmori 2022](#)).
- The US National Toxicology Program \$25 million animal study on long-term exposure to radiofrequency radiation found [DNA Damage, heart damage](#), increased [brain tumors, and increased heart tumors](#) deemed “clear evidence of cancer.” Importantly, this study was launched almost two decades ago by the FDA because the US government had not performed research on the long-term effects of RFR exposure and the FDA wanted data on long-term safety. In 1996, the EPA was defunded from developing proper safety standards, and since then there has been no systematic review of the science by any US agency.
- Researchers with the renowned Ramazzini Institute in Italy published [findings](#) that lab animals exposed to levels of RFR (below FCC limits) set to mimic cell tower exposures developed the same types of cancerous cancers as the [US National Toxicology Program](#) found in their large-scale animal study.
- In 2011, radiofrequency radiation was [classified](#) as a Class 2B possible carcinogen by the World Health Organization’s International Agency for Research on Cancer. This classification applies to RF regardless of the source. Between 2011 and today, the published peer-reviewed scientific evidence has significantly increased. Now, many scientists are of the opinion that the weight of current peer-reviewed evidence supports the conclusion that radiofrequency radiation should be regarded as a human carcinogen ([Hardell and Carlberg 2017](#), [Peleg et al, 2018](#), [Miller et al 2018](#)).
- An Australian [study](#) looked at RFR levels to which kindergarten children were exposed, depending on how close their school was to base stations/cell towers. Researchers equipped the children with RFR measuring devices. Researchers found that kindergartens located nearby base stations/cell towers (closer than 300 meters or approximately 330 yards) had total exposure to radiofrequency radiation (RFR or RF-EMF) more than 3 times higher than children at schools where base stations were further away than 300 meters.

- A 2018 [study](#) measured radiofrequency radiation exposures in the environment including emissions from cell phone towers, TV and FM radio broadcast antennas, cell phone handsets, and Wi-Fi—in several countries including the United States. The researchers concluded that cell phone tower (base station) radiation emissions are the dominant contributor to RFR exposure in most outdoor areas.
- A 2015 review found that in 93 out of 100 studies, RFR exposure caused oxidative stress ([Yakymenko 2015](#)). A 2021 review again confirmed non ionizing radiation has oxidative effects ([Schuermann 2021](#)). Many well-known causes of cancer in humans (such as asbestos and arsenic) are understood to induce oxidative stress.
- Studies also show that when combined with lead or a known carcinogen, RFR has magnified the carcinogen's effects. For example, RFR at levels far below FCC limits more than doubled the numbers of liver and lung tumors in carcinogen-exposed mice ([Lerchl 2015](#)).
- The International Association of Firefighters has officially opposed cell towers on their stations since 2004 after a study [found](#) neurological damage in firefighters with antennas on their fire station. In 2017, when 5G “small cells” were coming to California via a 5G streamlining bill (SB 649), firefighter organizations came out in strong opposition to the bill and requested that towers not be installed on firehouses. They were successful and SB649 was [amended](#) to [exempt](#) their stations from the deployment due to their health concerns.
- Published research finds the frequencies impact wildlife. For example, studies have found that the radiation alters bird navigation and disturbs honeybee colonies. Research also shows adverse impacts on trees and plants. ([Research on EMF and Bees](#), [Research on Wildlife](#) [Research on Trees](#))
- A 2019 [study](#) of students in schools near cell towers found their higher RF exposure was associated with impacts on motor skills, memory, and attention ([Meo 2019](#)). Examples of other effects linked to cell towers in research studies include [neuropsychiatric problems](#), [elevated diabetes](#), [headaches](#), [sleep problems](#), and [genetic damage](#). Such research continues to accumulate after the 2010 landmark [review study](#) on 56 studies that reported biological effects found at very low intensities of wireless radiation, including impacts on reproduction, permeability of the blood-brain barrier, behavior, cellular changes, and metabolic changes, and increases in cancer risk ([Lai and Levitt 2010](#)).
- The [International EMF Scientist Appeal](#) was submitted to the United Nations urging immediate protective policy action in light of the scientific evidence that has found adverse biological effects from electromagnetic radiation, including radiofrequency radiation, and, as of January 2019, this Appeal is signed by 247 scientists from 42 nations; these are scientists who have published peer-reviewed articles about electromagnetic fields. They state, “numerous recent scientific publications have shown that EMF affects living organisms at levels well below most international and national guidelines. Effects include increased cancer risk, cellular stress, increase in harmful free radicals, genetic damages, structural and functional changes of the

reproductive system, learning and memory deficits, neurological disorders, and negative impacts on general well-being.”

Please note that in several countries, governments have set policies to protect children, pregnant women, and medically fragile persons by classifying areas with homes, hospitals, and schools as “sensitive areas.” Some examples include:

- India lowered their RF limits to 1/10th of US FCC ICNIRP limits. *The Brihanmumbai Municipal Corporation, Zilla Parishad, Rajasthan, and Mumbai have banned cell antenna/tower installations on schools.*
- Greece has banned the installation of mobile phone base stations at the premises of schools, kindergartens, hospitals, or eldercare facilities.
- Chile’s “Antenna Law” prohibits cell antennas/towers in “sensitive areas” (educational institutions, nurseries, kindergartens, hospitals, clinics, nursing homes).
- Several countries have [lower allowable RFR limits](#) for “sensitive” areas which is generally defined as areas where children play and school.

EHT’s position is that children require special protections from radiofrequency radiation and their exposures should be reduced to as low as possible. We strongly recommend against cell tower/antenna placements at or near schools and homes as this would increase daily RFR exposure.

Please feel free to contact us with more questions.

Sincerely,

Devra Davis, PhD, MPH
President and Founder, Environmental Health Trust
Visiting Professor, Hebrew University Hadassah Medical Center
<https://ehtrust.org>

Anthony B. Miller, MD
Professor Emeritus at the Dalla Lana School of Public Health, University of Toronto
Senior Advisor to Environmental Health Trust

Theodora Scarato MSW Executive Director
Environmental Health Trust
Theodora.scarato@ehtrust.org

References

- Abdel-Rassoul, G., El-Fateh, O. A., Salem, M. A., Michael, A., Farahat, F., El-Batanouny, M., & Salem, E. (2007). Neurobehavioral effects among inhabitants around mobile phone base stations. *NeuroToxicology*, 28(2), 434–440. <https://doi.org/10.1016/j.neuro.2006.07.012>.
- Amraee, A., Seif, F., Bayatiani, M. R., Shakeri, M., & Zakeri, F. (2021). Correlation between Base Transceiver Station and the Quality of Sleep and Life of Nearby Residents. *Iranian Journal of Medical Physics*, 18(1), 10–14. https://ijmp.mums.ac.ir/article_14561.html.
- Balmori, A. (2022). Evidence for a health risk by RF on humans living around mobile phone base stations: From radiofrequency sickness to cancer. *Environmental Research*, 214, 113851. <https://doi.org/10.1016/j.envres.2022.113851>
- Balmori, A. (2010). Mobile phone mast effects on common frog (*Rana temporaria*) tadpoles: The city turned into a laboratory. *Electromagnetic Biology and Medicine*, 29(1–2), 31–35. <https://doi.org/10.3109/15368371003685363>.
- Blettner, M., Schlehofer, B., Breckenkamp, J., Kowall, B., Schmiedel, S., Reis, U., Potthoff, P., Schüz, J., & Berg-Beckhoff, G. (2009). Mobile phone base stations and adverse health effects: Phase 1 of a population-based, cross-sectional study in Germany. *Occupational and Environmental Medicine*, 66(2), 118–123. <https://doi.org/10.1136/oem.2007.037721>.
- Bortkiewicz, A., Zmyślony, M., Szykowska, A., & Gadzicka, E. (2004). [Subjective symptoms reported by people living in the vicinity of cellular phone base stations: Review]. *Medycyna Pracy*, 55(4), 345–351. <https://pubmed.ncbi.nlm.nih.gov/15620045/>.
- Broom, K. A., Findlay, R., Addison, D. S., Goiceanu, C., & Sienkiewicz, Z. (2019). Early-Life Exposure to Pulsed LTE Radiofrequency Fields Causes Persistent Changes in Activity and Behavior in C57BL/6 J Mice. *Bioelectromagnetics*, 40(7), 498–511. <https://doi.org/10.1002/bem.22217>
- Buchner, K., & Eger, H. D. I. (2011). *Changes of Clinically Important Neurotransmitters under the Influence of Modulated RF Fields A Long-term Study under Real-life Conditions*. <https://www.avaate.org/IMG/pdf/Rimbach-Study-20112.pdf>.
- Carlberg, M., Hedendahl, L., Koppel, T., & Hardell, L. (2019). High ambient radiofrequency radiation in Stockholm city, Sweden. *Oncology Letters*, 17(2), 1777–1783. <https://doi.org/10.3892/ol.2018.9789>.
- Choi, J., Min, K., Jeon, S., Kim, N., Pack, J.-K., & Song, K. (2020). Continuous Exposure to 1.7 GHz LTE Electromagnetic Fields Increases Intracellular Reactive Oxygen Species to Decrease Human Cell Proliferation and Induce Senescence. *Scientific Reports*, 10(1), 9238. <https://doi.org/10.1038/s41598-020-65732-4>
- Dode, A. C., Leão, M. M. D., Tejo, F. de A. F., Gomes, A. C. R., Dode, D. C., Dode, M. C., Moreira, C. W., Condessa, V. A., Albinatti, C., & Caiaffa, W. T. (2011). Mortality by neoplasia and cellular telephone base stations in the Belo Horizonte municipality, Minas Gerais state, Brazil. *The Science of the Total Environment*, 409(19), 3649–3665. <https://doi.org/10.1016/j.scitotenv.2011.05.051>.

Eger, et al., The Influence of Being Physically Near to a Cell Phone Transmission Mast on the Incidence of Cancer (2004). Umwelt·Medizin·Gesellschaft. <http://www.tetrawatch.net/papers/naila.pdf>.

Eskander, E. F., Estefan, S. F., & Abd-Rabou, A. A. (2012). How does long term exposure to base stations and mobile phones affect human hormone profiles? *Clinical Biochemistry*, 45(1–2), 157–161. <https://doi.org/10.1016/j.clinbiochem.2011.11.006>.

Eşmekaya, M. A., Seyhan, N., & Ömeroğlu, S. (2010). Pulse modulated 900 MHz radiation induces hypothyroidism and apoptosis in thyroid cells: A light, electron microscopy and immunohistochemical study. *International Journal of Radiation Biology*, 86(12), 1106–1116. <https://doi.org/10.3109/09553002.2010.502960>.

Falcioni, L., Bua, L., Tibaldi, E., Lauriola, M., De Angelis, L., Gnudi, F., Mandrioli, D., Manservigi, M., Manservigi, F., Manzoli, I., Menghetti, I., Montella, R., Panzacchi, S., Sgargi, D., Strollo, V., Vornoli, A., & Belpoggi, F. (2018). Report of final results regarding brain and heart tumors in Sprague-Dawley rats exposed from prenatal life until natural death to mobile phone radiofrequency field representative of a 1.8 GHz GSM base station environmental emission. *Environmental Research*, 165, 496–503. <https://doi.org/10.1016/j.envres.2018.01.037>

Gandhi, G., Kaur, G., & Nisar, U. (2015). A cross-sectional case control study on genetic damage in individuals residing in the vicinity of a mobile phone base station. *Electromagnetic Biology and Medicine*, 34(4), 344–354. <https://doi.org/10.3109/15368378.2014.933349>.

Gandhi, G., Naru, J., Kaur, M., & Kaur, G. (2014). DNA and Chromosomal Damage in Residents Near a Mobile Phone Base Station. *International Journal of Human Genetics*, 14(3–4), 107–118. <https://doi.org/10.1080/09723757.2014.11886234>.

Gómez-Perretta, C., Navarro, E. A., Segura, J., & Portolés, M. (2013). Subjective symptoms related to GSM radiation from mobile phone base stations: A cross-sectional study. *BMJ Open*, 3(12), e003836. <https://doi.org/10.1136/bmjopen-2013-003836>.

Hardell, L., & Koppel, T. (2022). Electromagnetic hypersensitivity close to mobile phone base stations—A case study in Stockholm, Sweden. *Reviews on Environmental Health*. <https://doi.org/10.1515/reveh-2021-0169>.

Hardell, L., Carlberg, M., Hedendahl, L. K., Koppel, T., & Ahonen, M. (2019). Environmental radiofrequency radiation at the Järntorget Square in Stockholm Old Town, Sweden in May, 2018 compared with results on brain and heart tumour risks in rats exposed to 1.8 GHz base station environmental emissions. *World Academy of Sciences Journal*, 1(1), 47–54. <https://doi.org/10.3892/wasj.2018.5>.

Hardell, L., Carlberg, M., & Hedendahl, L. K. (2018). Radiofrequency radiation from nearby base stations gives high levels in an apartment in Stockholm, Sweden: A case report. *Oncology Letters*, 15(5), 7871–7883. <https://doi.org/10.3892/ol.2018.8285>.

Hardell, L., Carlberg, M., Koppel, T., & Hedendahl, L. (2017). High radiofrequency radiation at Stockholm Old Town: An exposimeter study including the Royal Castle, Supreme Court, three major squares and the Swedish Parliament. *Molecular and Clinical Oncology*, 6(4), 462–476. <https://doi.org/10.3892/mco.2017.1180>.

Hardell, L., Koppel, T., Carlberg, M., Ahonen, M., & Hedendahl, L. (2016). Radiofrequency radiation at Stockholm Central Railway Station in Sweden and some medical aspects on public exposure to RF fields. *International Journal of Oncology*, 49(4), 1315–1324. <https://doi.org/10.3892/ijo.2016.3657>.

Hardell, L., & Sage, C. (2008). Biological effects from electromagnetic field exposure and public exposure standards. *Biomedicine & Pharmacotherapy*, 62(2), 104–109. <https://doi.org/10.1016/j.biopha.2007.12.004>.

Hecht, K., Savoley, E.N., (2007). Overloading of Towns and Cities with Radio Transmitters (Cellular Transmitter): a hazard for the human health and a disturbance of eco-ethics, IRCHET – International Research Centre of Healthy and Ecological Technology, Berlin, Germany. <https://ecfsapi.fcc.gov/file/7521097890.pdf>.

Hutter, H.-P., Moshhammer, H., Wallner, P., & Kundi, M. (2006). Subjective symptoms, sleeping problems, and cognitive performance in subjects living near mobile phone base stations. *Occupational and Environmental Medicine*, 63(5), 307–313. <https://doi.org/10.1136/oem.2005.020784>.

Khurana, V. G., Hardell, L., Everaert, J., Bortkiewicz, A., Carlberg, M., & Ahonen, M. (2010). Epidemiological evidence for a health risk from mobile phone base stations. *International Journal of Occupational and Environmental Health*, 16(3), 263–267. <https://doi.org/10.1179/107735210799160192>.

Koppel, T., & Hardell, L. (2022). Measurements of radiofrequency electromagnetic fields, including 5G, in the city of Columbia, SC, USA. *World Academy of Sciences Journal*, 4(3), 1–12. <https://doi.org/10.3892/wasj.2022.157>

Koppel, T., Ahonen, M., Carlberg, M., & Hardell, L. (2022). Very high radiofrequency radiation at Skeppsbron in Stockholm, Sweden from mobile phone base station antennas positioned close to pedestrians' heads. *Environmental Research*, 208, 112627. <https://doi.org/10.1016/j.envres.2021.112627>.

Koppel, T., Ahonen, M., Carlberg, M., Hedendahl, L. K., & Hardell, L. (2019). Radiofrequency radiation from nearby mobile phone base stations-a case comparison of one low and one high exposure apartment. *Oncology Letters*, 18(5), 5383–5391. <https://doi.org/10.3892/ol.2019.10899>.

Kundi, M., & Hutter, H.-P. (2009). Mobile phone base stations-Effects on wellbeing and health. *Pathophysiology: The Official Journal of the International Society for Pathophysiology*, 16(2–3), 123–135. <https://doi.org/10.1016/j.pathophys.2009.01.008>.

Levitt, B., & Lai, H. (2010). Biological effects from exposure to electromagnetic radiation emitted by cell tower base stations and other antenna arrays. *Environmental Reviews*, 18, 369–395. <https://doi.org/10.1139/a10-903>.

López, I., Félix, N., Rivera, M., Alonso, A., & Maestú, C. (2021). What is the radiation before 5G? A correlation study between measurements in situ and in real time and epidemiological indicators in Vallecas, Madrid. *Environmental Research*, 194, 110734. <https://doi.org/10.1016/j.envres.2021.110734>.

Lv, B., Chen, Z., Wu, T., Shao, Q., Yan, D., Ma, L., Lu, K., & Xie, Y. (2014). The alteration of spontaneous low frequency oscillations caused by acute electromagnetic fields exposure. *Clinical Neurophysiology*, 125(2), 277–286. <https://doi.org/10.1016/j.clinph.2013.07.018>

Marinescu, I. E., & Poparlan, C. (2016). Assessment of GSM HF-Radiation Impact Levels within the Residential Area of Craiova City. *Procedia Environmental Sciences*, 32, 177–183.
<https://doi.org/10.1016/j.proenv.2016.03.022>.

Meo, S. A., Almahmoud, M., Alsultan, Q., Alotaibi, N., Alnajashi, I., & Hajjar, W. M. (2019). Mobile Phone Base Station Tower Settings Adjacent to School Buildings: Impact on Students' Cognitive Health. *American Journal of Men's Health*, 13(1), 1557988318816914.
<https://doi.org/10.1177/1557988318816914>.

Meo, S. A., Alsubaie, Y., Almubarak, Z., Almutawa, H., AlQasem, Y., & Hasanato, R. M. (2015). Association of Exposure to Radio-Frequency Electromagnetic Field Radiation (RF-EMFR) Generated by Mobile Phone Base Stations with Glycated Hemoglobin (HbA1c) and Risk of Type 2 Diabetes Mellitus. *International journal of environmental research and public health*, 12(11), 14519–14528.
<https://doi.org/10.3390/ijerph121114519>.

Miller, A. B., Morgan, L. L., Udasin, I., & Davis, D. L. (2018). Cancer epidemiology update, following the 2011 IARC evaluation of radiofrequency electromagnetic fields (Monograph 102). *Environmental Research*, 167, 673–683. <https://doi.org/10.1016/j.envres.2018.06.043>

Navarro, E. A., Segura, J., Portolés, M., & Gómez-Perretta de Mateo, C. (2003). The Microwave Syndrome: A Preliminary Study in Spain. *Electromagnetic Biology and Medicine*, 22(2–3), 161–169.
<https://doi.org/10.1081/JBC-120024625>.

Oberfeld, G., Navarro, E., Portoles, M., Maestu, C., & Gómez-Perretta, C. (2002). *THE MICROWAVE SYNDROME - FURTHER ASPECTS OF A SPANISH STUDY*.
https://www.researchgate.net/publication/237410769_THE_MICROWAVE_SYNDROME_-_FURTHER_ASPECTS_OF_A_SPANISH_STUDY.

Özdemir, E., Çömelekoğlu, Ü., Degirmenci, E., Bayrak, G., Yildirim, M., Ergenoglu, T., Coşkun Yılmaz, B., Korunur Engiz, B., Yalin, S., Koyuncu, D. D., & Ozbay, E. (2021). The effect of 4.5 G (LTE Advanced-Pro network) mobile phone radiation on the optic nerve. *Cutaneous and Ocular Toxicology*, 40(3), 198–206. <https://doi.org/10.1080/15569527.2021.1895825>

Pachua, Lalrinthara & Pachua, Zaithanzauva. (2016). Health Effects of Mobile Tower Radiation on Human — Case Study. *International Journal of Applied Physics and Mathematics*. 6. 72-79.
[10.17706/ijapm.2016.6.2.72-79](https://doi.org/10.17706/ijapm.2016.6.2.72-79).

Pearce, J. M. (2020). Limiting liability with positioning to minimize negative health effects of cellular phone towers. *Environmental Research*, 181, 108845. <https://doi.org/10.1016/j.envres.2019.108845>.

Richter, E. D., Berman, T., & Levy, O. (2002). Brain cancer with induction periods of less than 10 years in young military radar workers. *Archives of Environmental Health*, 57(4), 270–272.
<https://doi.org/10.1080/00039890209601409>.

Roda, C., & Perry, S. (2014). Mobile phone infrastructure regulation in Europe: Scientific challenges and human rights protection. *Environmental Science & Policy*, 37, 204–214.
<https://doi.org/10.1016/j.envsci.2013.09.009>.

Rodrigues, N. C. P., Dode, A. C., de Noronha Andrade, M. K., O'Dwyer, G., Monteiro, D. L. M., Reis, I. N. C., Rodrigues, R. P., Frossard, V. C., & Lino, V. T. S. (2021). The Effect of Continuous Low-Intensity

Exposure to Electromagnetic Fields from Radio Base Stations to Cancer Mortality in Brazil. *International Journal of Environmental Research and Public Health*, 18(3), 1229.

<https://doi.org/10.3390/ijerph18031229>.

SA, M., Alsubaie, Y., Almubarak, Z., Almutawa, H., AlQasem, Y., & Hasanato, R. (2015). Association of Exposure to Radio-Frequency Electromagnetic Field Radiation (RF-EMFR) Generated by Mobile Phone Base Stations with Glycated Hemoglobin (HbA1c) and Risk of Type 2 Diabetes Mellitus. *International Journal of Environmental Research and Public Health*, 12, 14519-14528;

<https://doi.org/10.3390/ijerph121114519>.

Santini, R., Santini, P., Le Ruz, P., Danze, J. M., & Seigne, M. (2003). Survey Study of People Living in the Vicinity of Cellular Phone Base Stations. *Electromagnetic Biology and Medicine*, 22(1), 41–49.

<https://doi.org/10.1081/JBC-120020353>.

Santini, R., Santini, P., Danze, J. M., Le Ruz, P., & Seigne, M. (2002). Investigation on the health of people living near mobile telephone relay stations: I/Incidence according to distance and sex.

Pathologie-Biologie, 50(6), 369–373. [https://doi.org/10.1016/s0369-8114\(02\)00311-5](https://doi.org/10.1016/s0369-8114(02)00311-5). [Article in French].

Shinjo, T. & Shinjo, A. (2014) Significant Decrease of Clinical Symptoms after Mobile Phone Base Station Removal – An Intervention Study, Tetsuharu Shinjo and Akemi Shinjo
UmweltMedizinGesellschaft, 27(4), S. 294301.

Souffi, S., Lameth, J., Gaucher, Q., Arnaud-Cormos, D., Lévêque, P., Edeline, J.-M., & Mallat, M. (2022). Exposure to 1800 MHz LTE electromagnetic fields under proinflammatory conditions decreases the response strength and increases the acoustic threshold of auditory cortical neurons. *Scientific Reports*, 12(1), 4063. <https://doi.org/10.1038/s41598-022-07923-9>

Study of Cell Tower Radiation and its Health Hazards on human body - Lalrinthara Pachuau and Zaithanzauva Pachuau – IOSR Journal of Applied Physics (IOSR-JAP) e-ISSN: 2278-4861. Volume 6, Issue 1 Ver. 1 (Jan 2014), PP 01-06.

Vecsei, Z., Knakker, B., Juhász, P., Thuróczy, G., Trunk, A., & Hernádi, I. (2018). Short-term radiofrequency exposure from new generation mobile phones reduces EEG alpha power with no effects on cognitive performance. *Scientific Reports*, 8, 18010. <https://doi.org/10.1038/s41598-018-36353-9>

Wei, Y., Yang, J., Chen, Z., Wu, T., & Lv, B. (2019). Modulation of resting-state brain functional connectivity by exposure to acute fourth-generation long-term evolution electromagnetic field: An fMRI study. *Bioelectromagnetics*, 40(1), 42–51. <https://doi.org/10.1002/bem.22165>

Wolf, R., & Wolf, D. (2004). Increased incidence of cancer near a cell-phone transmitter station. *International Journal of Cancer*, 1(2), 123–128. [[Google Scholar](#)].

Yakymenko, I., Sidorik, E., Kyrilenko, S., & Chekhun, V. (2011). Long-term exposure to microwave radiation provokes cancer growth: Evidences from radars and mobile communication systems. *Experimental Oncology*, 33(2), 62–70. <https://pubmed.ncbi.nlm.nih.gov/21716201/>.

Yang, L., Chen, Q., Lv, B., & Wu, T. (2017). Long-Term Evolution Electromagnetic Fields Exposure Modulates the Resting State EEG on Alpha and Beta Bands. *Clinical EEG and Neuroscience*, 48(3), 168–175. <https://doi.org/10.1177/1550059416644887>

Yu, G., Tang, Z., Chen, H., Chen, Z., Wang, L., Cao, H., Wang, G., Xing, J., Shen, H., Cheng, Q., Li, D., Wang, G., Xiang, Y., Guan, Y., Zhu, Y., Liu, Z., & Bai, Z. (2020). Long-term exposure to 4G smartphone radiofrequency electromagnetic radiation diminished male reproductive potential by directly disrupting Spock3–MMP2–BTB axis in the testes of adult rats. *Science of The Total Environment*, 698, 133860. <https://doi.org/10.1016/j.scitotenv.2019.133860>

Zothansiam, Zosangzuali, M., Lalramdinpui, M., & Jagetia, G. C. (2017). Impact of radiofrequency radiation on DNA damage and antioxidants in peripheral blood lymphocytes of humans residing in the vicinity of mobile phone base stations. *Electromagnetic Biology and Medicine*, 36(3), 295–305. <https://doi.org/10.1080/15368378.2017.1350584>.